**Studies on chemical constituents of three
marine sponges, *Siphonochalina siphonella* in
Egypt and *Arenosclera* sp. and *Gelliodes* sp.
in Vietnam**

Dae-Won Ki

SECTION OF NATURAL PRODUCTS & DRUG DISCOVERY

INSTITUTE OF NATURAL MEDICINE

UNIVERSITY OF TOYAMA

2020

Table of Contents

Abbreviations.....	1
Chapter 1 Introduction	3
Chapter 2 Constituents of the <i>Siphonochalina siphonella</i> collected in Egypt and their cytotoxic activities	11
2.1 Introduction.....	12
2.2 Extraction and isolation	13
2.3 Structure elucidation of new compounds.....	16
2.3.1 Siphonellamide A (1)	16
2.3.2 Siphonellamide B (2)	21
2.3.3 Siphonellamide C (3)	26
2.3.4 Siphonellamide D (4)	29
2.3.5 Siphonellamide E (5)	33
2.3.6 Siphonellanol A (6)	37
2.3.7 Siphonellanol B (7)	42
2.3.8 Siphonellanol C (8).....	47
2.4 Cytotoxic activities of isolated compounds	51
2.5 The structure activity relationship (SAR) on cytotoxic activities.....	53
2.6 Summary of chapter 2.....	54
Chapter 3 Constituents of the <i>Arenosclera</i> sp. collected in Vietnam and their cytotoxic activities.....	55
3.1 Introduction.....	56
3.2 Extraction and isolation	57
3.3 Structure elucidation of new compounds.....	61
3.3.1 Tribromiododiphenyl ether (13).....	61
3.4 Cytotoxic activities of isolated compounds	66
3.5 The structure activity relationship (SAR) on cytotoxic activities.....	68
3.6 Antibacterial activities of isolated compounds	69
3.7 Summary of chapter 3	71
Chapter 4 Constituents of the <i>Gelliodes</i> sp. collected in Vietnam and their cytotoxic activities.....	72
4.1 Introduction.....	73
4.2 Extraction and isolation	74
4.3 Structure elucidation of new compounds.....	77
4.3.1 Gelliodesinic acid (26).....	77
4.4 Cytotoxic activities of isolated compounds	82

4.5	Summary of chapter 4	84
Discussion		85
Conclusion		86
Experimental		87
I.	General experimental procedures.....	87
II.	Experimental detail of chapter 2	89
III.	Experimental detail of chapter 3	95
IV.	Experimental detail of chapter 4	100
References		104
List of publications		116
Acknowledgements		117
Supplementary data		119

Abbreviations

$\mu\text{g/mL}$	Microgram per milliliter
μL	Micro liter
μM	Micro molar
A549	Human lung cancer cell line
^{13}C NMR	Carbon Nuclear Magnetic Resonance
^1H NMR	Proton Nuclear Magnetic Resonance
HOHAHA	Homonuclear Hartmann Hahn (TOCSY)
calcd	Calculated
CDCl_3	Deuterated chloroform
CH_3	Methyl
MeCN	Acetonitrile
CHCl_3	Chloroform
COSY	Correlation Spectroscopy
DMSO	Dimethyl sulfoxide
EtOAc	Ethyl acetate
g	Gram
HCl	Hydrochloric acid
HeLa	Human cervical cancer cell line
HMBC	Heteronuclear Multiple Bond Correlation
HMQC	Heteronuclear Multiple Quantum Correlation
HPLC	High Performance Liquid Chromatography
ESI-MS/MS	Electron Spray Ionization Mass Spectrometry MS/MS
HR-ESI-MS	High Resolution Electron Spray Ionization Mass Spectrometry
IC_{50}	Concentration of inhibitory causing 50% reduction in activity relative to the control
IR	Infrared
KBr	Potassium bromide
J	Coupling constant
m/z	Mass-to-charge ratio
MCF-7	Human breast cancer cell line

MeOH	Methanol
mg	Milligram
MHz	Megahertz
MIC	Minimum inhibitory concentration
mM	Millimolar
MS	Mass Spectroscopy
MTT	3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide
multi	Multiplicity
NMR	Nuclear Magnetic Resonance
NOESY	Nuclear Overhauser Effect Spectroscopy
NP	Normal Phase silica gel
PBDEs	Poly Brominated Diphenyl Ethers
TLC	Thin Layer Chromatography
TOCSY	Total Correlation Spectroscopy

Chapter 1 Introduction.

■ Introduction

Cancer has become the second leading cause of death worldwide. Globally, it is estimated that 1 in 6 deaths are caused by cancer and 9.6 million people died from cancer in 2018. It is reported that the most common form of cancer among male is lung cancer, which is account for 22% of all cancer cases. On the other hand, the highest mortality for the female is caused by breast cancer and cervical cancer, which account for 15% and 7.5%, respectively ¹⁻³. For many years, chemotherapeutic agents have been developed and used to treat cancer. However, there is no drug that shows good selectivity for cancer cells.

To date, about 70% of anti-cancer drugs were developed from natural resources, and world populations still rely mainly on anti-cancer drugs derived from medicinal plants ⁴. For examples, taxol has been isolated from the bark of the Western Yew tree, as useful drug for the treatment of breast cancer ⁵. Apart from taxol, vincristine and vinblastine isolated from Madagascar periwinkle plant have been served as an effective anti-cancer agent against childhood leukemia, Hodgkin's disease, and choriocarcinoma ⁶. Few marine anti-cancer agents have also reached to clinical phase trials. One example is bryostatin 1, which has been recently entered phase II clinical trial against melanoma, non-Hodgkin's lymphoma, renal cancer, and colorectal cancer ⁷⁻¹⁰. Furthermore, several marine natural products exhibited *in vitro* anti-cancer activity and cytotoxic activity against a wide range of the cancer cell lines. However, marine natural products have not been yet fully exploited ¹¹⁻¹³.

Among the marine organisms, marine sponges belong to phylum Porifera and comprises of approximately 8,000 different species in the world. These marine organisms attach themselves to solid surfaces such as rocks, shells or sea floor ^{14–16}. The shapes of marine sponges can range from the simplest tubular shape to complex features like the cup-shaped and treelike shape or even amorphous ¹⁷. Due to the sedentary life, marine sponges produce many bioactive secondary metabolites for defense purposes. The research of bioactive secondary metabolites from marine sponges started as early as the 1950s ^{18–21}. In the past decade, the studies on marine sponges have gained a lot of attention due to the discovery of new bioactive substances that are important to the pharmaceutical field ^{22–24}. This phenomenon has further accelerated, especially from the year 2001 to 2010. During these years, more than 200 new compounds per year have been discovered from marine sponges and marine sponges have been shown to be rich source of wide variety of chemically distinct bioactive compounds ^{17, 25–29}. The newly discovered compounds from marine sponges can be generally classified into 4 main classes, terpenoids, alkaloids, steroids, and peptides, and among them, terpenoids reportedly occupied about 26.0% followed by alkaloids (22.5%), steroids (9.5%), and peptides (7.8%) (Figure 1.1) ^{17, 25–27, 30–36}. Remarkably, almost half (49.2%) of the isolated compounds have exhibited cytotoxic activity against cancer cell lines, suggesting high potential of the compounds in marine sponges as cancer drug seeds (Figure 1.2) ¹⁷.

Figure 1.1 Chemical classes of new compounds isolated from marine sponges ¹⁷

Figure 1.2 Isolation of new bioactive compounds from marine sponge ¹⁷

Furthermore, more than 50% of all new cytotoxic compounds isolated from marine sponges between 2001 to 2010 were discovered from the orders Dictyoceratida, Poecilosclerida, and Haplosclerida of marine sponges; Dictyoceratida, Poecilosclerida, and Haplosclerida have provided 22.0%, 17.3 %, and 12.1 of all cytotoxic compounds,

respectively. In particular, marine sponges belonging to the Dysideidae, Irciniidae, Spongiidae, and Thorectidae families in Dictyoceratida have provided numerous structurally diverse new cytotoxic compounds, such as trisnorsesterterpenoids, thiopyrone, iodo-sesquiterpene, sesquiterpene benzoxazole, furanoterpenoids, manoalide, and alkaloids (Figure 1.3) ^{37–42}. The marine sponges of Acarnidae, Coelosphaeridae, Crambeidae, Hymedesmiidae, Latrunculiidae, and Podospongiidae families containing in Poecilosclerida have provided vast cytotoxic bispyrroloiminoquinone, sesterterpenoids, norterpenoids, and alkaloids (Figure 1.3). The Haplosclerida marine sponge has also been known to be the rich source of acetylenic carotenoids, polyacetylene diol, alkylpyridine, linear acetylene, cyclic peptides, and glycerol ether lipids with cytotoxic activities (Figure 1.3) ^{43–46}.

Figure 1.3 New cytotoxic compounds from marine sponge A. New cytotoxic compounds from marine sponge Dictyoceratida, B. New cytotoxic compounds from marine sponge Poecilosclerida, C. New cytotoxic compounds from marine sponge Haplosclerida

Countries with long coast line such as Egypt and Vietnam have abundant marine resources. However, the marine resources from these countries were less exploited, especially for the discovery of new cytotoxic compounds. Hence, in this study, in order

to search for new cytotoxic compounds effective to three most common cancers in female and male, a marine sponge, *Siphonochalina siphonella* (Haplosclerida), and seven marine sponges, *Callyspongia* sp. (Haplosclerida), *Iotrochota* sp. (Poecilosclerida), *Gelliodes* sp. (Haplosclerida), *Hyrtilia* sp. (Dictyoceratida), *Petrosia* sp. (Haplosclerida), *Arenosclera* sp. (Haplosclerida), and *Gelliodes fibulata* (Haplosclerida) were collected from Egypt and Vietnam, respectively, and the MeOH extracts from the marine sponges were screened for cytotoxic activities against human cervical cancer cell line (HeLa), human breast cancer cell line (MCF-7), and human lung cancer cell line (A549). The screening revealed that the MeOH extracts of *S. siphonella*, *Arenosclera* sp., and *Gelliodes* sp. showed cytotoxic activities against HeLa, MCF-7, and A549 with IC₅₀ values of each cell line 25.5, 7.9, and 36.7 µg/mL, 13.0, 8.5, and 25.4 µg/mL, 27.5, 9.0, and 32.2 µg/mL, respectively (Table 1.1). *S. siphonella*, *Arenosclera* sp., and *Gelliodes* sp. are most likely to be the potential source for discovery of cytotoxic compounds.

Table 1.1 Cytotoxic activities of marine sponge extract

Samples	IC ₅₀ (µg/ml)		
	HeLa	MCF-7	A549
<i>S. siphonella</i>	25.5	13.0	27.5
<i>Callyspongia</i> sp.	> 100	> 100	> 100
<i>Iotrochota</i> sp.	> 100	> 100	> 100
<i>Geolliodes</i> sp.	36.7	25.4	32.2
<i>Hyrtilia</i> sp.	90.4	82.6	> 100
<i>Petrosia</i> sp.	> 100	91.7	> 100
<i>Arenosclera</i> sp.	7.9	8.5	9.0
<i>G. fibulata</i>	> 100	> 100	> 100

Thus, this study aimed to obtain new cytotoxic compounds from *S. siphonella* in Egypt and *Arenosclera* sp. and *Gelliodes* sp. in Vietnam, and further investigation of the

extract from these marine sponges led to isolation of ten new compounds and 21 known compounds.

Chapter 2 will discuss about the isolation and structure elucidation of eight new polyacetylene derivatives from *S. siphonella* and their cytotoxic activities. The isolation and structural elucidation of one new polybrominated iododiphenyl ether from *Arenosoclea* sp. as well as cytotoxic activities of the isolated compounds will be reported in chapter 3. Finally, isolation and structure elucidation of new compounds from the marine sponge *Gelliodes* sp. and their cytotoxic activities will be discussed in chapter 4.

Chapter 2 Constituents of *Siphonochalina siphonella* collected in Egypt and their cytotoxic activities

2.1 Introduction

S. siphonella (*Callyspongia siphonella*) belongs to the Callyspongiidae family. *S. siphonella* consists of several tubes and ending in a central oval shaped osculums (26–40 μm diameter) and are colored in red, pink, or lavender ^{51, 52}. Sponges from this genus is endemic to the Red Sea.

Studies had been conducted on *S. siphonella* to isolate compounds with biological activities. One of the most prevalent compounds are triterpenoid such as sipholane, neviotane, siphonellane, and dahabane ^{53–55}. These compounds were reported to possess anti-osteoclastogenesis ⁵⁶, anti-cancer ^{57, 58}, anti-bacteria ⁵⁹, anti-viral ⁵⁹, and cytotoxicity ^{59, 60}. Polyacetylene compounds were also isolated from *S. siphonella* ^{61, 62}. However, these compounds could be also found in other marine sponge species. As an example, *C. fistularis* was reported to produce a compound with a rare structure, callyspongamide A, which consists of a phenyl moiety attached to a linear polyacetylene chain structure ⁶³. Polyacetylene compounds were also reported to possess potent cytotoxic activities against cancer cell lines, a human breast cancer cell line MCF-7 ⁶¹, human cervical cancer cell line HeLa ⁶³, murine leukemia cancer cell line P388 ⁶⁴, human skin cancer cell line A2058 ⁶⁵, and human lung cancer cell lines H522-T1 and H460 ⁶⁵.

In agreement with previous studies, the MeOH extract of *S. siphonella* exhibited cytotoxic activities against HeLa, MCF-7, and A549 (Table 1.1). Hence, further study has been performed to investigate new chemical constituents with cytotoxic activities from Egyptian marine sponge *S. siphonella*.

2.2 Extraction and isolation

S. siphonella. (600 g wet weight) was macerated in MeOH (2 L $\times$ 3) for 2 h with sonication. The extract was evaporated under vacuum, and the remaining aqueous residue was successively partitioned with CHCl₃ (4.5 g). A CHCl₃ fraction was subjected to normal phase silica gel open column chromatography using solvent systems composed of *n*-hexane/EtOAc and EtOAc/MeOH with increasing polarities. In total, 190 fractions of 30 mL each were collected and pooled into 21 combined fractions after TLC profiling. These fractions were further subjected to chromatographic separation and preparative HPLC (Figure 2.2), which led to isolation of twelve compounds including three new polyacetylenic amides, siphonellamide A (**1**), siphonellamide B (**2**), and siphonellamide C (**3**), a new monoacetylene amide, siphonellamide D (**4**), a new fatty indole amide siphonellamide E (**5**), and three new polyacetylenic alcohols, siphonellanol A (**6**), siphonellanol B (**7**), and siphonellanol C (**8**), together with four known compounds, *N*-[2''-(1'*H*-indol-3'-yl)ethyl] -hexadecanamide (**9**)⁶⁶, callyspongamide A (**10**)⁶³, dehydroisophonochalynol (**11**)⁵¹, and siphonochalynol (**12**)⁵¹ (Figure 2.3).

Figure 2.1 *Siphonochalina siphonella*

Figure 2.2 Extraction and isolation of compounds from MeOH extract of *S. siphonella*

Figure 2.3 Structures of isolated compounds 1–12 from *S. siphonella*

2.3 Structure elucidation of new compounds

2.3.1 Siphonellamide A (1)

Compound **1** was obtained as a yellowish oil. The positive HR-ESI-MS showed a pseudo-molecular ion peak at m/z 414.2759 $[M+Na]^+$ (calcd. for $C_{27}H_{37}NONa$, 414.2773) and molecular formula was established as $C_{27}H_{37}NO$ in conjugation with NMR data (Table 2.1). The UV spectrum exhibited maximum absorptions at 211, 215, 218, and 228 nm. The IR spectrum showed absorption bands at 3434 cm^{-1} (N–H), 3287 cm^{-1} (C≡H), 2926 cm^{-1} , 2854 cm^{-1} (C–H), 2352 cm^{-1} (C≡C), and 1648 cm^{-1} (C=C). The 1H NMR spectrum of **1** revealed the five aromatic proton signals at δ_H 7.32 (2H, t, $J = 7.6$ Hz, H-5' and H-7'), 7.23 (tt, $J = 7.6, 1.4$ Hz, H-6'), and 7.20 (2H, dd, $J = 7.6, 1.4$ Hz, H-4' and H-6'), two olefinic methine signals at δ_H 5.99 (dt, $J = 10.9, 7.5$ Hz, H-16) and 5.43 (d, $J = 10.9$ Hz, H-17), a terminal acetylene signal at δ_H 3.07 (d, $J = 2.1$ Hz, H-19), and 14 methylenes, which accounted for the remaining protons in the molecules (Figure 2.4 and Table 2.1). The ^{13}C NMR and HMQC spectra exhibited a carbonyl carbon signal at δ_C 171.5 (C-1), an aromatic quaternary carbon signal at δ_C 138.9 (C-3'), three aromatic methine carbon signals at δ_C 128.8 (overlapped, C-4' and C-8'), 128.6 (overlapped, C-5' and C-7'), and 126.5 (C-6'), two mono substituted olefinic carbon signals at δ_C 146.2 (C-16) and 107.9 (C-17), and four acetylenic carbons including three non-protonated carbon signals at δ_C 81.2 (overlapped, C-5), 80.7 (C-18), and 79.5 (C-4) and a protonated carbon signal at δ_C 81.2 (overlapped, C-19), together with 14 methylene carbons (Figure 2.5 and Table 2.1). These NMR data of **1** were similar to callyspongamide A, which was isolated by Youssef et al (2003)⁶³ from *Callyspongia fistularis*. The significant difference between **1** and callyspongamide A was the abduction of the two methylene units in **1**.

The 1H – 1H COSY correlations of C(4')H–C(5')H–C(6')H–C(7')H–C(8')H and

$C(1')H_2-C(2')H_2$ as well as the HMBC correlations from $H-5'$ to $C-3$, from $H-7'$ to $C-3'$, from H_2-2' to $C-1'$, $C-4'$, and $C-8'$, and from H_2-1' to $C-3'$ established the presence of an ethyl benzene moiety in **1** (Figure 2.6). The ESI-MS/MS experiments measured on the pseudo-molecular ion at m/z 431 $[M+H]^+$ showed an ion fragment at m/z 105 $[M-C_{19}H_{28}NO]^+$, which further confirmed the presence of ethyl benzene moiety in **1**. The HMBC correlations from H_2-1' to $C-1$, from H_2-3 to $C-1$, from H_2-2 to $C-4$, and from H_2-7 to $C-5$, the $^1H-^1H$ COSY correlation between H_2-6 and H_2-7 , and the HOHAHA long range correlation of H_2-3 to H_2-7 indicated the presence of a hept-4-ynamide moiety ($C-1-C-6$) in **1** (Figure 2.6). One of the most important key HMBC correlation of H_2-1' to $C-1$ suggested that the ethyl benzene moiety connected to the hept-4-ynamide moiety in **1** (Figure 2.6). The ESI-MS/MS ion fragment at m/z 200 $[M-C_{14}H_{23}]^+$ supported this connection of **1**. Meanwhile, the HOHAHA long range correlation of $H-17$ to $H-19$ and the $^1H-^1H$ COSY correlations of $C(14)H_2-C(15)H_2-C(16)H-C(17)H$ established the (3*Z*)-pent-3-en-1-yne moiety. The ESI-MS/MS ion fragments at m/z 121 $[M+H-C_{19}H_{27}O]^+$ and 200 $[M-C_{14}H_{23}]^+$ and HR-ESI-MS deduced the presence of an aliphatic moiety located from $C-8$ to $C-13$. The key COSY correlation of $C(15)H_2-C(14)H_2$ showed a connection of the aliphatic and (3*Z*)-pent-3-en-1-yne moieties (Figure 2.6). Compound **1** was thus determined to be a (16*Z*)-*N*-(2-phenylethyl)nonadec-16-en-4,18-diynamide, and was named siphonellamide A.

Figure 2.4 ^1H NMR (400 MHz) spectrum of siphonellamide A (**1**)

Figure 2.5 ^{13}C NMR (100 MHz) spectrum of siphonellamide A (**1**)

Figure 2.6 Key HMBC (arrows), ^1H - ^1H COSY (bold lines), and HOHAHA (dotted lines) correlations of **1**

Figure 2.7 ESI-MS/MS fragmentation patterns of **1**

Table 2.1 ^1H (400 MHz) and ^{13}C NMR (100 MHz) data for **1** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	1	
	δ_{H} (J in Hz)	δ_{C}
1		171.5
2	2.15 (m)	36.2
3	2.29 (m)	18.6
4		79.5
5		81.2*
6	2.13 (m)	18.5
7	1.47 (m)	28.7
8	1.30* (m)	28.8
9	1.30* (m)	29.3
10	1.25* (m)	29.0
11	1.25* (m)	28.8
12	1.30* (m)	28.6
13	1.30* (m)	29.1*
14	1.49 (m)	29.1*
15	2.32 (m)	30.2
16	5.99 (dt, 10.9, 7.5)	146.2
17	5.43 (d, 10.9)	107.9
18		80.7
19	3.07 (d, 2.1)	81.2*
1'	3.53 (td, 6.9, 6.0)	40.5
2'	2.82 (t, 6.9)	35.7
3'		138.9
4'	7.20 (dd, 7.6, 1.4)	128.8*
5'	7.32 (t, 7.6)	128.6*
6'	7.23 (tt, 7.6, 1.4)	126.5
7'	7.32 (t, 7.6)	128.6*
8'	7.20 (dd, 7.6, 1.4)	128.8*

* Overlapping resonances within the same column.

2.3.2 Siphonellamide B (2)

Compound **2** was obtained as a yellowish oil. The HR-ESI-MS displayed a pseudo-molecular ion peak at m/z 453.2865 $[M+Na]^+$ (calcd. for $C_{29}H_{38}N_2ONa$, 453.2882) and the molecular formula of **2** was established as $C_{29}H_{38}N_2O$ in conjugation with the NMR data (Table 2.2). The UV spectrum exhibited maximum absorptions at 218, 227, and 290 nm. The IR spectrum showed absorption bands at 3422 cm^{-1} (N–H), 3276 cm^{-1} (C≡H), 2926 cm^{-1} , 2860 cm^{-1} (C–H), 2356 cm^{-1} (C≡C), 1728 cm^{-1} (C=O), and 1642 cm^{-1} (C=C).

The 1H and ^{13}C NMR spectra in conjugation of HMQC spectrum indicated the presence of an indole unit in **2**, [δ_H 7.05 (s, H-2')/ δ_C 122.3(C-2'), δ_H 7.39 (dd, $J = 8.1, 1.0$ Hz, H-4')/ δ_C 111.2 (C-4'), δ_H 7.14 (ddd, $J = 8.1, 7.0, 1.0$ Hz, H-5')/ δ_C 119.5 (C-5'), δ_H 7.22 (ddd, $J = 7.9, 7.0, 1.0$ Hz, H-6')/ δ_C 122.0(C-6'), δ_H 7.62 (d, $J = 7.9$ Hz, H-7')/ δ_C 118.7 (C-7'), δ_C 136.6 (C-7'a), 127.2(C-3'a), 113.1 (C-3')], instead of the phenyl moiety in **1**. These NMR data suggested that **2** possesses an indole moiety instead of phenyl moiety. The spin systems $[C(4')H-C(5')H-C(6')H-(7')H]$ and $[NH-C(1'')H_2-C(2'')H_2]$ in the 1H - 1H COSY spectrum and cross peaks from H-7' and H-6' to C-7'a, from H-5' and H-4' to C-3'a, from H-2' to C-3'a and C-7'a, and from H-2'' to C-1'', C-2', C-3', and C-3'a in the HMBC spectrum confirmed the presence of the ethyl indole moiety in **2** (Figure 2.10), which was further evident from ESI-MS/MS ion fragments at m/z 144 $[M-C_{19}H_{30}NO]^+$, 160 $[M+H-C_{19}H_{27}O]^+$, and 187 $[M-C_{18}H_{27}]^+$ (Figure 2.11). In addition, the HMQC spectrum also revealed the presence of an amide carbonyl [δ_C 172.7 (C-1)], a non-protonated acetylene unit [δ_C 81.2 (C-5) and 79.6 (C-4)], a terminal acetylene unit [δ_H 3.07 (d, $J = 2.3$ Hz, H-19)/ δ_C 81.2 (C-19) and δ_C 80.6 (C-18)], and a *cis*-oriented olefin

unit [δ_{H} 6.00 (dt, $J = 10.7, 7.0$ Hz, H-16)/ δ_{C} 146.3 (C-16) and 5.44 (d, $J = 10.7$ Hz, H-17)/ δ_{C} 107.9 (C-17)]. Comparisons of the NMR data and HR-ESI-MS data of **2** with those of **1** deduced the presence of a nonadeca-16-en-4,18-diynamide moiety in **2**. The HMBC correlations from H₂-2 to C-1 and C-4 and from H₂-7 to C-5, the ¹H–¹H COSY correlations [C(14)H₂–C(15)H₂–C(16)H₂–C(17)H₂], and the HOHAHA long range correlation of H-17 to H-19 supported the presence of the moiety in **2**. The HMBC correlation from H₂-1'' to C-1 confirmed a connection between indole and (Z)-nonadeca-16-en-4,18-diynamide moieties. Compound **2** was determined to be (16Z)-N-[2-(1H-indol-3-yl)ethyl]nonadec-16-en-4,18-diynamide, and was named siphonellamide B.

Figure 2.8 ^1H NMR (400 MHz) spectrum of siphonellamide B (**2**)

Figure 2.9 ^{13}C NMR (100 MHz) spectrum of siphonellamide B (**2**)

Figure 2.10 Key HMBC (arrows), ^1H - ^1H COSY (bold lines), and HOHAHA (dotted lines) correlations of **2**

Figure 2.11 ESI-MS/MS fragmentation patterns of **2**

Table 2.2. ^1H (400 MHz) and ^{13}C NMR (100MHz) data for **2** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	2	
	δ_{H} (J in Hz)	δ_{C}
1		172.7
2	2.11 (m)	36.3
3	2.13* (m)	18.5
4		79.6
5		81.2*
6	2.13* (m)	18.7
7	1.48 (m)	28.7
8	1.30* (m)	29.1*
9	1.30* (m)	29.3
10	1.30* (m)	28.6
11	1.30* (m)	29.1*
12	1.30* (m)	29.0*
13	1.30* (m)	29.0*
14	1.46 (m)	28.8
15	2.33 (m)	30.2
16	6.00 (dt, 10.7, 7.0)	146.3
17	5.44 (d, 10.7)	107.9
18		80.6
19	3.07 (d, 2.3)	81.2*
-NH	5.51 (br s)	
1''	3.61 (td, 6.7, 6.0)	39.6
2''	2.99 (t, 6.7)	25.4
-NH	8.08 (br s)	
2'	7.05 (s)	122.3
3'		113.1
3'a		127.2
4'	7.39 (dd, 8.1, 1.0)	111.2
5'	7.14 (ddd, 8.1, 7.0, 1.0)	119.5
6'	7.22 (ddd, 7.9, 7.0, 1.0)	122.0
7'	7.62 (d, 7.9)	118.7
7'a		136.6

* Overlapping resonances within the same column.

2.3.3 Siphonellamide C (**3**)

Compound **3** was obtained as a yellowish oil. The HR-ESI-MS of **3** displayed a pseudo-molecular ion peak at m/z 451.2712 $[M+Na]^+$ (calcd. for $C_{29}H_{36}N_2ONa$, 451.2725). The molecular formula of **3** revealed an additional degree of unsaturation as compared with that of **2**. The carbon signals of **3** were assigned by the HMQC and HMBC cross peaks. The 1H NMR spectrum (Figures 2.12 and Table 2.3), in association with the HMQC data of **3** were quite similar to those of **2**. The significant difference between **3** and **2** was the olefinic proton regions, where δ_H 5.82 (dt, $J = 10.8, 7.4$ Hz, H-12) and δ_H 5.42 (br d, $J = 10.8$ Hz, H-13) were exhibited in **3**, suggesting that **3** has two *cis*-oriented olefin units.

The 1H - 1H COSY correlations $[C(10)H_2-C(11)H_2-C(12)H-C(13)H]$ and the HOHAHA long range correlation of H_2-10 to $H-16$ presented a *cis*-oriented olefin unit $[\delta_H$ 5.82 (dt, $J = 10.8, 7.4$ Hz, H-12)/ δ_C 143.2 (C-12) and δ_H 5.42 (br d, $J = 10.8$ Hz, H-13)/ δ_C 136.0 (C-13)] in **3**. The ESI-MS/MS fragment at m/z 105 $[M-C_{21}H_{27}N_2O]^+$ confirmed this olefin location (C-12–C-13) in **3**. The remaining partial structure was determined by comparison with NMR data of **2** and ESI-MS/MS fragment pattern analyses (Figures 2.13 and 2.14). Compound **3** was determined to be (12Z,16Z)-*N*-[2-(1*H*-indol-3-yl)ethyl]nonadec-12,16-dien-4,18-diynamide, and was named siphonellamide C.

Figure 2.12 ^1H NMR (500 MHz) spectrum of siphonellamide C (**3**)

Figure 2.13 Key HMBC (arrows), ^1H - ^1H COSY (bold lines), and HOHAHA (dotted lines) correlations of **3**

Figure 2.14 ESI-MS/MS fragmentation patterns of **3**

Table 2.3. ^1H (500 MHz) and ^{13}C NMR data by 2D NMR spectrum for **3** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	3	
	δ_{H} (J in Hz)	δ_{C}
1		173.3
2	2.14 (m)	36.0
3	2.32 (m)	18.8
4		79.0
5		82.8
6	2.34 (m)	19.2
7	1.53 (m)	28.6
8	1.54 (m)	29.6
9	1.39 (m)	29.8*
10	1.40 (m)	29.8*
11	2.27 (m)	30.6*
12	5.82 (dt, 10.8, 7.4)	143.2
13	5.42 (br d, 10.8)	136.0
14	1.73 (m)	25.0
15	2.31 (m)	30.6*
16	5.99 (dt, 10.8, 7.0)	146.3
17	5.44 (d, 10.8)	110.0
18		80.7
19	3.07 (d, 2.3)	81.3
-NH	5.50 (br s)	
1''	3.63 (td, 6.7, 6.0)	40.4
2''	2.99 (t, 6.7)	25.9
-NH	8.04 (br s)	
2'	7.05 (s)	122.6
3'		113.3
3'a		127.3
4'	7.39 (dd, 8.0, 1.0)	111.5
5'	7.14 (ddd, 8.0, 7.0, 1.0)	119.1
6'	7.22 (ddd, 8.0, 7.0, 1.0)	122.4
7'	7.62 (dd, 8.0, 1.0)	118.8
7'a		136.5

2.3.4 Siphonellamide D (**4**)

Compound **4** was obtained as a yellowish oil. The HR-ESI-MS of **4** displayed a pseudo-molecular ion peak at m/z 441.2881 $[M+Na]^+$ (calcd. for $C_{28}H_{38}N_2ONa$, 441.2882). The IR and UV-spectra of **4** exhibited similar absorption patterns to **2** and **3**. The 1H NMR spectrum of **4** showed similarity to acetylene amide type compound (Table 2.4). In the molecular formula, the significant differences in **4** were the presence of two protons and absence of a carbon, suggesting that **4** lacks the acetylene moiety (C-4–C-5) of **3**. The 1H NMR and 2D NMR spectra of **4** exhibited the signals corresponding to an indole unit, a terminal acetylene unit [δ_H 3.07 (d, $J = 2.3$ Hz, H-18)/ δ_C 81.3 (C-18) and δ_C 80.7 (C-17)], an amide carbonyl unit [δ_C 173.6 (C-1)], two *cis*-oriented olefin units [δ_H 5.34 (dt, $J = 10.8, 6.4$ Hz, H-4)/ δ_C 130.6 (C-4), δ_H 5.33 (dt, $J = 10.8, 6.4$ Hz, H-5)/ δ_C 130.6 (C-5), δ_H 5.99 (dt, $J = 10.9, 7.5$ Hz, H-15)/ δ_C 146.0 (C-15), δ_H 5.44 (d, $J = 10.9$ Hz, H-16)/ δ_C 108.3 (C-16)], and 13 methylenes (Figures 2.15 and Table. 2.4).

The HMBC correlations from H-1'' to C-1/C-3' and from H-2'' to C-1''/C-3' and the 1H – 1H COSY correlations [NH–C(1'')H₂–C(2'')H₂] indicated the presence of *N*-2-(1*H*-indol-3-yl)ethyl amide moiety. The spin systems [C(14)H₂–C(15)H–C(16)H] in the 1H – 1H COSY spectrum and the long range correlation of H-16/H-18 in the 1H – 1H TOCSY spectrum confirmed the presence of a (3*Z*)-pent-3-en-1-yne moiety. The fragment structure (C-2–C-6) was determined from 2D NMR data. The 1H – 1H COSY correlations [C(2)H₂–C(3)H₂–C(4)H–C(5)H–C(6)H₂] and the HMBC correlations from H-2 to C-1 and from H-7 to C-5 indicated the presence of *cis*-oriented olefin unit [δ_H 5.34 (dt, $J = 10.8, 6.4$ Hz, H-4)/ δ_C 130.6 (C-4), δ_H 5.33 (dt, $J = 10.8, 6.4$ Hz, H-5)/ δ_C 130.6 (C-5)] at C-4 and C-5 of **4** (Figures 2.16). A ESI-MS/MS ion fragment at m/z 163 [M-

$C_{16}H_{19}N_2O]^+$ supported this olefin location in **4** (Figure 2.17). This ESI-MS/MS data confirmed that the seven methylenes located between C-6 and C-14. Compound **4** was determined to be (4*Z*,15*Z*)-*N*-[2-(1*H*-indol-3-yl)ethyl]octadec-4,15-dien-17-ynamide and was named siphonellamide D.

Figure 2.15 ^1H NMR (500 MHz) spectrum of siphonellamide D (**4**)

Figure 2.16 Key HMBC (arrows), ^1H - ^1H COSY (bold lines), and HOHAHA (dotted lines) correlations of **4**

Figure 2.17 ESI-MS/MS fragmentation patterns of **4**

Table 2.4. ^1H (500 MHz) and ^{13}C NMR data by 2D NMR spectrum for **4** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	4	
	δ_{H} (J in Hz)	δ_{C}
1		173.6
2	2.10 (m)	36.8
3	2.02 (m)	27.1
4	5.34 (dt, 10.8, 6.4)	130.6*
5	5.33 (dt, 10.8, 6.4)	130.6*
6	2.00 (m)	27.1
7	1.41 (m)	31.8
8	1.30* (m)	29.3*
9	1.30* (m)	29.3*
10	1.26* (m)	29.7*
11	1.26* (m)	29.7*
12	1.30* (m)	29.3*
13	1.30* (m)	29.3*
14	2.32 (m)	30.1
15	5.99 (dt, 10.9, 7.5)	146.0
16	5.44 (d, 10.9)	108.3
17		80.7
18	3.07 (d, 2.3)	81.3
-NH	5.47 (br s)	
1''	3.62 (td, 6.7, 6.0)	40.7
2''	2.98 (t, 6.7)	25.4
-NH	8.02 (br s)	
2'	7.05 (s)	122.4
3'		113.2
3'a		127.7
4'	7.39 (dd, 8.1, 1.0)	111.1
5'	7.14 (ddd, 8.1, 7.0 1.0)	119.7
6'	7.23 (ddd, 8.0, 7.0, 1.0)	122.9
7'	7.62 (dd, 8.0, 1.0)	118.8
7'a		136.6

* Overlapping resonances within the same column.

2.3.5 Siphonellamide E (**5**)

Compound **5** was obtained as a yellowish oil. The HR-ESI-MS displayed a pseudo-molecular ion peak at m/z 419.3025 $[M+Na]^+$ (calcd. for $C_{26}H_{40}N_2ONa$, 419.3038) and the molecular formula of **5** was established as $C_{26}H_{40}N_2O$ in conjugation with the NMR data (Table 2.5). The UV and IR spectra were similar to compounds **2–4**. The 1D-NMR spectra of **5** displayed typical signals, corresponding to an indole unit, two olefin carbons at δ_C 129.8 and 130.0, and a carbonyl carbon at δ_C 173.2. However, the NMR data of **5** didn't show acetylene moiety signals (Figure 2.18, 2.19 and Table 2.5). These NMR data of **5** were nearly identical to those of the known compound **9** that has been isolated from *Pseudomonas* spp.⁶⁶. Only the significant difference in **5** was a olefinic signals [δ_H 5.35 (m, H-8)/ δ_C 129.8 (C-8) and δ_H 5.35 (m, H-9)/ δ_C 130.0 (C-9)].

The location of the olefin was confirmed to be between C-8 and C-9 by ozonolysis, which showed a ESI-MS ion fragment at m/z 301 $[M+H]^+$, consistent with a *N*-[2-(1*H*-indol-3-yl)-ethyl]-8-oxooctanamide unit (Figure S34). The geometric configuration of the Δ^8 -double bond was identified as *cis* configuration, based on the δ_C 27.2 ppm for C-7 and C-10^{49, 67}. Other partial structure of **5** was determined by comparison of NMR data of **2–4**. Compound **5** was identified as (8*Z*)-*N*-[2-(1*H*-indol-3-yl)ethyl] hexadec-8-enamide and was named siphonellamide E.

Figure 2.18 ^1H NMR (400 MHz) spectrum of siphonellamide E (5)

Figure 2.19 ^{13}C NMR (100 MHz) spectrum of siphonellamide E (5)

Figure 2.20 Key HMBC (arrows) and ¹H-¹H COSY (bold lines) correlations of **5**

Figure 2.21 ESI-MS/MS fragmentation patterns of **5**

Table 2.5. ^1H (400 MHz) and ^{13}C NMR (100 MHz) data of **5** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	5	
	δ_{H} (J in Hz)	δ_{C}
1		173.2
2	2.09 (m)	36.9
3	1.57 (m)	25.3
4	1.25* (m)	29.0
5	1.25* (m)	29.2
6	1.25* (m)	29.3
7	2.01* (m)	27.2*
8	5.35* (m)	129.8
9	5.35* (m)	130.0
10	2.01* (m)	27.2*
11	1.25* (m)	29.6*
12	1.25* (m)	29.6*
13	1.25* (m)	27.2*
14	1.25* (m)	31.8
15	1.25* (m)	22.6
16	0.87 (t, 6.9)	14.1
-NH	5.50 (br s)	
1''	3.62 (td, 6.7, 6.0)	39.6
2''	2.99 (t, 6.7)	25.3
-NH	8.10 (br s)	
2'	7.04 (s)	122.2
3'		113.1
3'a		127.3
4'	7.39 (dd, 8.1, 1.0)	111.2
5'	7.13 (ddd, 8.1, 7.0, 1.0)	119.5
6'	7.21 (ddd, 8.1, 7.0, 1.0)	122.0
7'	7.62 (d, 8.1)	118.7
7'a		136.4

* Overlapping resonances within the same column.

2.3.6 Siphonellanol A (**6**)

Compound **6** was obtained as a yellowish oil. The HR-ESI-MS displayed a pseudo-molecular ion peak at m/z 325.1543 $[M+Na]^+$ (calcd. for $C_{22}H_{22}ONa$, 325.1568) and the molecular formula of **6** was established as $C_{22}H_{22}O$ in conjugation with the NMR data (Table 2.6). This molecular formula indicated 12 indices of hydrogen deficiency. The UV spectrum exhibited maximum absorptions at 213, 218, and 229 nm. The IR spectrum showed absorption bands at 3432 cm^{-1} (O–H), 3303 cm^{-1} ($C\equiv H$), 2928 cm^{-1} , 2865 cm^{-1} (C–H), 2355 cm^{-1} , 2213 cm^{-1} ($C\equiv C$), and 1639 cm^{-1} ($C=C$). The 1D-NMR and HMQC spectra of **6** exhibited signals of a terminal acetylene unit [δ_H 3.09 (d, $J = 1.8\text{ Hz}$, H-22)/ δ_C 81.8 (C-22) and δ_C 80.5 (C-21)], a *cis*-oriented olefin unit [δ_H 5.98 (dt, $J = 10.8, 7.4\text{ Hz}$, H-19)/ δ_C 145.3 (C-19), δ_H 5.48 (d, $J = 10.8\text{ Hz}$, H-20)/ δ_C 108.9 (C-20)], a *trans*-oriented olefin unit [δ_H 5.54 (d, $J = 15.8\text{ Hz}$, H-4)/ δ_C 110.2 (C-4), δ_H 6.11 (dt, $J = 15.8, 7.8\text{ Hz}$, H-5)/ δ_C 143.6 (C-5)], an oxygenated methylene unit [δ_H 4.37 (br s, H-1)/ δ_C 51.7 (C-1)], eight non-protonated acetylene carbon signals at δ_C 86.4 (C-2), 84.2 (C-3), 79.2 (C-9), 78.4 (C-14), 66.5 (C-13), 66.0 (C-11), 60.3 (C-12), and 60.0 (C-11), and eight methylene signals (Figure 2.22, 2.23, and Table 2.6). The NMR data of **6** were similar to those of dehydroisophonochalynol (**11**) isolated from the same sponge⁵¹ as well as in this study, except for the coupling constant at H-4 and H-5. The coupling constants at H-4 and H-5 in **6** showed 15.8 Hz, while those of **11** were observed as 11.0 Hz. It suggested that **6** has a *trans*-oriented olefin unit in C-4–C-5 instead of the *cis*-oriented olefin of **11** (Table 2.6).

The 1H – 1H COSY correlation [$C(4)H$ – $C(5)H$ – $C(6)H_2$] and the HMBC correlations from H_2 -1 to C-2, from H-5 to C-3, and from H_2 -6 to C-4 and C-5 established

a (*E*)-hept-4-en-2-yn-1-ol moiety (C-1–C-7). The long range correlations of H₂-1 to H₂-6 and H₂-7 to H₂-1/H-4/H-5 in the ¹H–¹H TOCSY spectrum supported the presence of this subunit. The spin systems [C(18)H₂–C(19)H–C(20)H] in the ¹H–¹H COSY spectrum, the cross peaks from H₂-16 to C-17 and from H₂-18 to C-19/C-20 in the HMBC spectrum, and long range correlations of H₂-16 to H-19/H-20/H-22 in ¹H–¹H TOCSY spectrum indicated the presence of a (*Z*)-hept-3-en-1-yne (C-16–C-22) in **6**. The molecular formula of **6** and comparisons of NMR data of **6** and **11** suggested that **6** has an octa-2,4,6-triyne unit (C-8–C-15) as same as **11**. The HMBC correlation from H₂-7 to C-9 allowed connection between the (*E*)-hept-4-en-2-yn-1-ol and octa-2,4,6-triyne units (C-8–C-15). Therefore, the subunit (*Z*)-hept-3-en-1-yne (C-16–C-22) connected to the opposite side of the octa-2,4,6-triyne unit (C-8–C-15). The ESI-MS/MS experiment measured on the pseudo-molecular ion peak at *m/z* 325 [M+Na]⁺ displayed ion fragments at *m/z* 180 [M+H–C₈H₁₁O]⁺, 208 [M+H–C₆H₇O]⁺, and 237 [M–C₅H₅]⁺ (Figure 2.24). These fragmentation patterns supported the assignments in **6**. The compound **6** was determined to be (4*E*,19*Z*)-docosa-4,19-dien-2,9,11,13,21-pentayn-1-ol and was named siphonellanol A.

Figure 2.22 ^1H NMR (500 MHz) spectrum of siphonellanol A (6)

Figure 2.23 ^{13}C NMR (125 MHz) spectrum of siphonellanol A (6)

Figure 2.24 Key HMBC (arrows), ^1H - ^1H COSY (bold lines), and HOHAHA (dotted lines) correlations of **6**

Figure 2.25 ESI-MS/MS fragmentation patterns of **6**

Table 2.6. ^1H and ^{13}C NMR data of **6** and **11** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	6^a		11^b	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
1	4.37 (br s)	51.7	4.40 (d, 2.0)	51.4
2		86.4		92.0
3		84.2		81.8
4	5.54 (d, 15.8)	110.2	5.48 (d, 11.0)	109.8
5	6.11 (dt, 15.8, 7.8)	143.6	5.83 (dt, 11.0, 7.3)	142.4
6	2.24 (m)	31.9	2.35 (m)	29.0
7	1.64 (m)	27.2	1.60 (quint, 7.3)	27.1
8	2.31* (m)	18.7	2.27* (m)	18.7
9		79.2		79.1
10		66.0		65.7
11		60.0		60.2
12		60.3		60.4
13		66.5		65.9
14		78.4		78.8
15	2.31* (m)	19.4	2.27* (m)	19.0
16	1.54 (m)	27.5	1.50 (m)	27.3
17	1.52 (m)	27.8	1.51 (m)	27.6
18	2.33 (m)	29.6	2.28 (m)	29.4
19	5.98 (dt, 10.8, 7.4)	145.3	5.94 (dt, 10.9, 7.8)	145.0
20	5.48 (d, 10.8)	108.9	5.42 (d, 10.9)	108.5
21		80.5		80.2
22	3.09 (d, 1.8)	81.8	3.04 (d, 1.8)	81.5

^a ^1H NMR and ^{13}C NMR were measured at 500 MHz and 125 MHz, respectively.

^b ^1H NMR and ^{13}C NMR were measured at 400 MHz and 100 MHz, respectively.

* Overlapping resonances within the same column.

2.3.7 Siphonellanol B (7)

Compound **7** was obtained as a yellowish oil. The HR-ESI-MS of **7** displayed the pseudo-molecular ion peak at m/z 329.1858 $[M+Na]^+$ (calcd. for $C_{22}H_{26}ONa$, 329.1881). The IR and UV spectrum exhibited similar absorption pattern to **6**. The 1D-NMR data, in conjunction with the HMQC experiment of **7**, showed the signals corresponding to a (*Z*)-pentadeca-3-en-1,9,11,13-tetrayne unit (C-8–C-22) as same as **6** (Table 2.7, Figure 2.26 and 2.27). The assignment was confirmed by 1H – 1H COSY, HMBC, 1H – 1H TOCSY, and ESI-MS/MS fragmentation analyses. The 1H – 1H COSY correlations $[C(17)H_2-C(18)H_2-C(19)H-C(20)H]$, HMBC correlations from H_2 -16 to C-14/C-17 and H_2 -18 to C-19/C-20, HOHAHA long range correlations of H_2 -17 to H-19/H-20 and H-20 to H-22, and ESI-MS/MS ion fragment at m/z 180 $[M+H-C_8H_{11}O]^+$ confirmed the presence of the (*Z*)-pentadeca-3-en-1,9,11,13-tetrayne unit. Meanwhile, the 1D-NMR data of **7** revealed three methylene signals including an oxygenated methylene $[\delta_H$ 3.67 (t, J = 6.5 Hz, H-1)/ δ_C 62.6 (C-1), δ_H 1.61 (m, H-2)/ δ_C 32.6 (C-2), δ_H 2.32 (m, H-3)/ δ_C 29.7 (C-3)]. These NMR data suggested that **7** lacks the non-protonated acetylene moiety observed in **6**. In addition, the NMR spectrum also showed a *cis*-oriented olefin unit $[\delta_H$ 5.34 (dt, J = 11.0, 7.3 Hz, H-4)/ δ_C 130.5 (C-4), δ_H 5.44 (dt, J = 11.0, 7.2 Hz, H-5)/ δ_C 128.8 (C-5)]. The 1H – 1H COSY spectrum of **7** showed two spin systems, $[C(1)H_2-C(2)H_2-C(3)H_2]$ and $[C(4)H-C(5)H-C(6)H_2-C(7)H_2]$. The HMBC correlations from H_2 -1 to C-2, from H_2 -3 to C-4/C-5, and from H_2 -7 to C-4 and HOHAHA long range correlations of H_2 -1 to H-5/ H_2 -6/ H_2 -7 confirmed the presence of the subunit (C-1–C-7) in **7**. The ESI-MS/MS ion fragments at m/z 261 $[M-C_2H_5O]^+$ and 180 $[M+H-C_8H_{15}O]^+$ supported this assignment in **7**. Therefore, **7** was determined to be (4*Z*,19*Z*)-

docosa-4,19-dien-9,11,13,21-tetrayne-1-ol and was named siphonellanol B.

Figure 2.26 ^1H NMR (500 MHz) spectrum of siphonellanol B (7)

Figure 2.27 ^{13}C NMR (100 MHz) spectrum of siphonellanol B (7)

Figure 2.28 Key HMBC (arrows), ^1H - ^1H COSY (bold lines), and HOHAHA (dotted lines) correlations of **7**

Figure 2.29 ESI-MS/MS fragmentation patterns of **7**

Table 2.7. ^1H (500 MHz) and ^{13}C NMR (100 MHz) data of **7** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	7	
	δ_{H}	δ_{C}
1	3.67 (t, 6.5)	62.6
2	1.61 (m)	32.6
3	2.32 (m)	29.7
4	5.34 (dt, 11.0, 7.3)	130.5
5	5.44 (dt, 11.0, 7.2)	128.8
6	2.13 (m)	26.0
7	1.56 (m)	27.4
8	2.30* (m)	18.8
9		72.9
10		65.9
11		60.4
12		60.4
13		65.9
14		79.0
15	2.30* (m)	19.2
16	1.54 (m)	27.9
17	1.52 (m)	27.6
18	2.37 (m)	29.5
19	5.98 (dt, 10.8, 7.5)	145.2
20	5.48 (d, 10.8)	108.7
21		80.3
22	3.09 (d, 1.8)	81.5

* Overlapping resonances within the same column.

2.3.8 Siphonellanol C (**8**)

Compound **8** was obtained as a colorless oil. The HR-ESI-MS displayed a pseudo-molecular ion peak at m/z 327.1708 $[M+Na]^+$ (calcd. for $C_{22}H_{24}ONa$, 327.1725) and the molecular formula of **8** was established as $C_{22}H_{24}O$, in conjugation with the NMR data. The IR and UV spectra of **8** exhibited similar absorption patterns to **6** and **7**. The HMQC data of **8** was similar to those of **7** (Figure 2.30, 2.31, and Table 2.8). However, the NMR data of **8** indicated the presence of a *trans*-oriented olefin unit [δ_H 6.41 (dd, J = 15.0, 10.7 Hz, H-5)/ δ_C 126.3 (C-5), δ_H 5.71 (dt, J = 15.0, 7.7 Hz, H-6)/ δ_C 131.2 (C-6)] and a *cis*-oriented olefin unit [δ_H 5.38 (dt, J = 10.7, 7.7 Hz, H-3)/ δ_C 127.0 (C-3), δ_H 6.12 (t, J = 10.7 Hz, H-4)/ δ_C 132.4 (C-4)]. The 1H - 1H COSY correlations, [C(1)H₂-C(2)H₂-C(3)H-C(4)H] and [C(5)H-C(6)H-C(7)H₂], the HMBC correlations from H₂-2 to C-1/C-3/C-4, and the HOHAHA long range correlations of H₂-7 to H₂-1/H-3/H-4/H-5 confirmed that these two olefins were located at C-5-C-6 and C-3-C-4, respectively. The ESI-MS/MS ion fragments, m/z 180 $[M+H-C_8H_{11}O]^+$ and 197 $[M-C_8H_{11}]^+$, supported the assignment of the olefin units in **8** (Figure 2.32 and 2.33). The partial structures (C-8-C-22) of **8** were determined by comparison of NMR data with those of **6** and **7** as well as 1H - 1H COSY, HMBC, and 1H - 1H TOCSY spectra of **8**. Therefore, **8** was identified to be (3*Z*,5*E*,19*Z*)-docosa-3,5,19-trien-9,11,13,21-tetrayne -1-ol and was named siphonellanol C.

Figure 2.30 ^1H NMR (400 MHz) spectrum of siphonellanol C (**8**)

Figure 2.31 ^{13}C NMR (100 MHz) spectrum of siphonellanol C (**8**)

Figure 2.32 Key HMBC (arrows), ^1H - ^1H COSY (bold lines), and HOHAHA (dotted lines) correlations of **8**

Figure 2.33 ESI-MS/MS fragmentation patterns of **8**

Table 2.8. ^1H (400 MHz) and ^{13}C NMR (100 MHz) data of **8** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	8	
	δ_{H}	δ_{C}
1	3.69 (t, 6.4)	62.2
2	2.45 (q, 6.4)	31.2
3	5.38 (dt, 10.7, 7.7)	127.0
4	6.12 (t, 10.7)	132.4
5	6.41 (dd, 15.0, 10.7)	126.3
6	5.71 (dt, 15.0, 7.7)	131.2
7	2.39 (m)	31.4
8	2.36* (m)	19.7
9		73.0
10		65.8
11		60.6
12		60.2
13		65.8
14		79.2
15	2.36* (m)	19.2
16	1.60* (m)	27.7
17	1.60* (m)	27.4
18	2.37 (m)	29.5
19	5.98 (dt, 11.0, 7.5)	145.2
20	5.42 (d, 11.0)	108.7
21		80.3
22	3.09 (d, 2.3)	81.5

* Overlapping resonances within the same column.

2.4 Cytotoxic activities of isolated compounds from *S. siphonella*

The cytotoxic activities of all isolated compounds (**1**, **2**, and **5–12**) were tested against human cervical cancer cell line HeLa, human breast cancer cell line MCF-7, and human lung cancer cell line A549 at the concentrations of 10, 50, and 100 μM , using MTT assay. Compounds **3** and **4** were not evaluated on cytotoxic activities due to their low yield. All of the tested compounds except for **5** and **9** exhibited cytotoxic activities against the three cancer cell lines. Compound **5** exhibited weak cytotoxic activity against HeLa, whereas this compound didn't show activity against MCF-7 and A549. Compounds **1** and **2** showed cytotoxic activities against the three cancer cell lines with IC_{50} values ranging from 9.4 μM to 34.1 μM . Compounds **6–8** showed moderate cytotoxic activities against HeLa with IC_{50} values of 26.5, 26.2, and 25.9 μM , respectively, while **6–8** exhibited moderate cytotoxic activities against MCF-7 and A549 with IC_{50} values ranging from 54.9 to 69.2 μM . The assay result agrees well with previous literature, where **10** possessed moderate cytotoxic activity against HeLa with the IC_{50} value of 25.5 μM ⁶³. In addition, **10** showed moderate cytotoxic activities against MCF-7 and A549 with IC_{50} values of 19.4 and 30.0 μM , respectively. The known compounds **11** and **12** displayed moderate cytotoxic activities against the three cancer cell lines with IC_{50} values ranging from 23.9 to 63.4 μM . The previous study also reported that **11** exhibited moderate activities against HeLa and MCF-7, which are similar to the assay results^{61, 64}.

Table 2.9. Cytotoxic activities of isolated compounds from *S. siphonella*

Samples	IC ₅₀ (μM)		
	HeLa	MCF-7	A549
1	9.4	18.0	24.2
2	17.4	34.1	25.9
5	78.4	> 100	> 100
6	26.5	54.9	59.8
7	26.2	69.2	59.9
8	25.9	57.6	58.5
9	> 100	> 100	> 100
10	25.5	19.4	30.0
11	23.9	61.4	63.4
12	25.4	61.3	60.4
5-FU ^a	28.4	34.7	14.6

^a 5-Fluorouracil: positive control.

2.5 The structure–activity relationship (SAR) on cytotoxic activities.

Comparisons of the cytotoxic results of the isolated compounds confirmed that the acetylene moiety plays an important role in the cytotoxic activities, as previously reported. Furthermore, comparisons of the structure and activity of phenyl polyacetylene amides **1** and **10**, indole polyacetylene amide **2**, and polyacetylene alcohols **6–8**, **11**, and **12** suggested that the combination of the acetylene moiety with the phenyl or indole moiety could increase cytotoxic activities against MCF-7 and A549.

However, in this study, the effects of the positions and number of acetylene moieties in the structures of the isolated polyacetylene compounds on cytotoxic activities still remained unclear.

2.6 Summary of chapter 2

Three new polyacetylene amides, siphonellamide A (**1**), siphonellamide B (**2**), and siphonellamide C (**3**), one new monoacetylene amide, siphonellamide D (**4**), one indole fatty amide, siphonellamide E (**5**), three new polyacetylene alcohols, siphonellanol A (**6**), siphonellanol B (**7**), and siphonellanol C (**8**), together with four known compounds (**9–12**) were isolated from the CHCl₃ fraction of the MeOH extract of the Egyptian marine sponge *S. siphonella*. Polyacetylene amides **1**, **2**, and **10** exhibited potent cytotoxic activities against the three tested cancer cell lines, HeLa, MCF-7, and A549, with the IC₅₀ values ranging from 9.4 to 34.1 µM. Meanwhile, polyacetylene alcohols **6–8**, **11**, and **12** displayed moderate cytotoxic activities with the IC₅₀ values ranging from 23.9 to 69.2 µM against HeLa, MCF-7, and A549. However, the indole fatty amides **5** and **9** didn't show cytotoxic activities against MCF-7 and A549 except for HeLa with the IC₅₀ value of 78.4 µM.

The MTT assay suggested that polyacetylene amides and polyacetylene alcohols possessed moderate cytotoxic activities against three tested cancer cell lines. Among them, polyacetylene amides showed stronger cytotoxic activity against MCF-7 and A549 than polyacetylene alcohols. Furthermore, these cytotoxic results suggested that the combination of acetylene moiety with indole moiety or phenyl moiety were crucial for increasing the cytotoxic activities against MCF-7 and A549.

Therefore, the isolation of indole fatty amides or polyacetylene amides from *S. siphonella* may provide a new chemotaxonomic incentive to this sponge and the biosynthetic origin of the polyacetylene amides may be due to cyanobacteria in *S. siphonella*.

Chapter 3 Constituents of *Arenosclera* sp. collected in Vietnam and their cytotoxic activities

3.1 Introduction

The genus *Arenosclera* belongs to the Callyspongiidae family. The sponge *Arenosclera* sp. are white, cream or beige in color, and exhibited soft texture mass ⁵². It has circular or oval-shaped osculans (9 mm in diameter) and spicules of up to 110 μ m in length ⁶⁸. These sponges are endemic to the Southwest Pacific (Australia), New Caledonia, and South China Sea ^{52, 69}.

To date, only *A. brasiliensis* from the *Arenosclera* genus yielded compounds with biological activities ^{70, 71}. The compounds isolated from *A. brasiliensis* were tetracyclic alkylpiperidine alkaloids, haliclonacyclamine E and arenosclerins A–C, which possess antimicrobial activities against *Staphylococcus aureus*, *Escherichia coli*, *P. aeruginosa*, and *C. albicans* and cytotoxic activities against human leukemia cell line HL-60, murine fibrosarcoma cancer cell line L929, human skin cancer cell line B16, and human brain cancer cell line U138 ^{70, 71}.

In the preliminary study, the MeOH extract of *Arenosclera* sp. harvested from Vietnam exhibited cytotoxic activities against HeLa, MCF-7, and A549. Therefore, further study was performed to investigate chemical constituents of *Arenosclera* sp. with new cytotoxic compounds.

3.2 Extraction and isolation

Arenosclera sp. (474 g dry weight) was macerated in MeOH (2 L $\times$ 3) for 2 h with sonication. The extract (79.7 g) was evaporated under vacuum and was subjected to HP-20 column chromatography using MeOH and H₂O to give H₂O, 50% MeOH (0.6 g), and MeOH (52.5 g) fractions, respectively. MeOH fraction was subjected to normal phase silica gel open column chromatography using solvent systems composed of *n*-hexane/EtOAc and EtOAc/MeOH with increasing polarities. In total, 83 fractions of 500 mL each was collected and pooled into 15 combined fractions after TLC profiling. These fractions were further separated by repeated column chromatography (Figure 3.2), which furnished one new compound together with eight known polybrominated diphenyl ethers (PBDEs) and three sterols. The new 3-bromo-2-(2',4'-dibromophenoxy)-6-iodophenol (**13**) was named tribromiododiphenyl ether. Eight known compounds were identified as 2-(2',4'-dibromophenoxy)-4,6-dibromophenol (**14**)⁷², 2-(4'-bromophenoxy)-3,5-dibromophenol (**15**)⁷³, 5-bromo-2-(2',4'-dibromophenoxy)phenol(**16**)⁷⁴, 2-(2',4'-dibromophenoxy)-5,6-dibromo-3-chlorophenol (**17**)⁷⁵, 2-(2',4'-dibromophenoxy)-3,5-dibromophenol (**18**)⁷⁵, 2-(2',4'-dibromophenoxy)-3,5,6-tribromophenol (**19**)⁷⁵, 3,6-dibromo-2-(2',4'-dibromophenoxy)phenol (**20**)⁷⁶, 2-(2',4'-dibromophenoxy)-4,6-dibromoanisole (**21**)⁷⁷, cholesterol (**22**)⁷⁸, 5 α , 8 α , epidioxysterol-6-en-3 β -ol (**23**)⁷⁹, and inseparable epidioxysterol (**24-a/b**)⁸⁰ by comparisons of NMR data with literatures previously reported those one (Figure 3.3).

Figure 3.1 *Arenosclera* sp.

Figure 3.2 Extraction and isolation of compounds from MeOH extract of *Arenosclera* sp.

Figure 3.3 Structures of isolated compounds **13**–**24** from *Arenoscelra* sp.

3.3 Structure elucidation of new compounds

3.3.1 Tribromoiiododiphenyl ether (**13**)

Compound **13** was obtained as a white powder and its HR-ESI-MS exhibited a pseudo molecular ion peak at m/z 550.6822 $[M+6-H]^-$, 548.6853 $[M+4-H]^-$, 546.6889 $[M+2-H]^-$, and 544.6859 $[M-H]^-$ (calcd. for $C_{12}H_5O_2Br_3I$: 544.6890), corresponding to the molecular formula of $C_{12}H_6O_2Br_3I$ in conjunction with NMR data. These isotope ion peaks indicated three bromines in **13**. The UV spectrum displayed maximum absorptions at 229, 283, and 290 nm. The IR spectrum showed absorption bands at 3428 cm^{-1} (phenolic O-H), 2920 cm^{-1} (aromatic C-H str), 1577 cm^{-1} (aromatic C=C str), 1465 cm^{-1} (aromatic C=C bend), 1093 cm^{-1} (C-O str), 690 cm^{-1} (C-Br), and 550 cm^{-1} (C-I). The 1H NMR spectrum of **13** revealed five aromatic protons at δ_H 7.78 (d, $J = 2.3$ Hz, H-3'), 7.51 (d, $J = 8.6$ Hz, H-5), 7.30 (dd, $J = 8.6, 2.3$ Hz, H-5'), 7.00 (d, $J = 8.6$ Hz, H-4), and 6.43 (d, $J = 8.6$ Hz, H-6') and a hydroxyl proton at δ_H 5.83 (br s, 1-OH). Five protonated benzene carbons at δ_C 126.5 (C-4), 136.3 (C-5), 136.1 (C-3'), 131.4 (C-5'), and 116.0 (C-6') and seven quaternary carbons at δ_C 149.8 (C-1), 138.7 (C-2), 117.4 (C-3), 82.9 (C-6), 152.3 (C-1'), 112.7 (C-2'), and 115.8 (C-4') were revealed from the ^{13}C NMR and HMQC spectra of **13** (Figure 3.4 and 3.5). These NMR data of **13** were similar to those of **20**, a polybrominated diphenyl ether (PBDE). The only difference between **13** and **20** was that **13** had an up-field shifted non-protonated benzene carbon at δ_C 82.9 (C-6), instead of δ_C 109.8 (C-6) in **20**. It suggested that **13** possesses an iodine atom at C-6.

The 1H - 1H COSY correlation of C(5')H-C(6')H, the HMBC correlations from H-3' to C-1', C-2', C-4', and C-5', from H-5' to C-1' and C-3', and from H-6' to C-1', C-2', and C-4' indicated the presence of a 2',4'-dibromophenoxy moiety in **13** (Figure 3.6).

Moreover, the spin system [C(4)H–C(5)H] in the ^1H – ^1H COSY spectrum and the cross peaks from H-4 to C-2, C-3, and C-6 and from H-5 to C-1, C-3, and C-6 in the HMBC spectrum established a 3-bromophenol moiety in **13** (Figure 3.6). These signals indicated the 2',4'-dibromophenoxy moiety attached to the 3-bromophenol moiety with an oxygen atom. The HR-ESI-MS ion fragments at m/z 311.1689 $[\text{M-H-C}_6\text{H}_3\text{OBr}_2]^-$ and 248.9590 $[\text{M-H-C}_6\text{H}_3\text{OBrI}]^-$ supposed the presence of the iodine in the 3-bromophenol moiety in **13** (Figure 3.7). Hence, **13** was determined to be a 3-bromo-2-(2',4'- dibromophenoxy)-6-iodophenol through the NMR spectra and was named tribromiododiphenyl ether.

Figure 3.4 ^1H NMR (500 MHz) spectrum of tribromoiododiphenylether (13)

Figure 3.5 ^{13}C NMR (125 MHz) spectrum of tribromoiododiphenylether (13)

Figure 3.6 HMBC (arrows) and ¹H-¹H COSY (bold lines) of tribromiodo-diphenylether (**13**)

Figure 3.7 HR-ESI-MS ion fragments of tribromiododiphenylether (**13**)

Table 3.1 ^1H (500 MHz) and ^{13}C NMR (125 MHz) data of **13** and **20** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	13		20	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
1		149.8		147.3
1-OH	5.83 (br. s)		5.84 (br. s)	
2		138.7		139.8
3		117.4		116.6
4	7.00 (d, 8.6)	126.5	7.12 (d, 8.6)	125.4
5	7.51 (d, 8.6)	136.3	7.31 (d, 8.6)	130.2
6		82.9		109.8
1'		152.3		152.4
2'		112.7		112.6
3'	7.78 (d, 2.3)	136.1	7.78 (d, 2.3)	136.0
4'		115.8		115.6
5'	7.30 (dd, 8.6, 2.3)	131.4	7.28 (dd, 8.6, 2.3)	131.3
6'	6.43 (d, 8.6)	116.0	6.42 (d, 8.6)	115.9

3.4 Cytotoxic activities of isolated compounds from *Arenosclera* sp.

All isolated compounds (**13–24**) were tested for cytotoxic activities against human cervical cancer cell line HeLa, human breast cancer cell line MCF-7, and human lung cancer cell line A549 at the concentrations of 10, 50, and 100 μM , using the MTT assay (Table. 3.2). Most of the isolated PBDEs, except for **18** and **20**, showed cytotoxic activities against the three cancer cell lines. The new compound **13** exhibited moderate cytotoxic activity against only MCF-7 with an IC_{50} value of 35.6 μM , but didn't show any activity against HeLa and A549. Compound **14** displayed strong cytotoxic activity against the three cancer cell lines, HeLa, MCF-7, and A549, with IC_{50} values ranging from 5.0 to 7.5 μM . Compound **15** showed strong cytotoxic activity against MCF-7 with an IC_{50} value of 5.1 μM and moderate cytotoxic activities against HeLa and A549 with IC_{50} values of 38.6 and 68.5 μM , respectively. Compound **16** displayed weak cytotoxic activity against HeLa with an IC_{50} value of 86.8 μM and moderate cytotoxic activity against MCF-7 with an IC_{50} value of 68.3 μM , but didn't show cytotoxic activity against A549. In addition, **17**, **19**, and **21** exhibited moderate cytotoxic activities against the three cancer cell lines with IC_{50} values ranging from 43.3 to 78.3 μM . The isolated three sterols didn't exhibit cytotoxic activities against the three cancer cell lines.

Table 3.2 Cytotoxic activities of isolated compounds from *Arenosclera* sp.

Samples	IC ₅₀ (μM)		
	HeLa	MCF-7	A549
13	> 100	35.6	> 100
14	6.9	5.0	7.5
15	38.6	5.1	68.5
16	86.8	68.3	> 100
17	72.3	64.1	73.2
18	> 100	> 100	> 100
19	62.0	43.3	78.3
20	> 100	> 100	> 100
21	65.7	51.4	39.7
22	> 100	> 100	> 100
23	> 100	> 100	> 100
24a/b	> 100	> 100	> 100
5-FU ^a	43.1	46.2	47.5

^a 5-Fluorouracil: positive control.

3.5 The structure–activity relationship (SAR) on cytotoxic activities.

Comparisons of **14**, **18**, and **20** suggested that the *para*-orientation relationship between the hydroxyl and bromine groups at C-1 and C-4 was the most important feature for enhancing the cytotoxic activities. On the other hands, in the case of the lack of the bromine moiety from C-4, as confirmed from the comparisons of **13**, **18**, **20**, and **21**, the presence of either iodine moiety at C-6 or the methoxy group at C-1 were important for increasing the activities.

3.6 Antibacterial activities of isolated compounds from *Arenosclera* sp.

All isolated compounds were also evaluated for their antibacterial activities against gram positive bacteria, *B. subtilis* and *S. aureus*, and gram negative bacteria, *K. pneumonia* and *E. coli*. Compound **13** exhibited strong antibacterial activities against *B. subtilis*, *S. aureus*, and *K. pneumoniae*, with MIC values ranging between 3.1–6.3 μ M. Compound **14** showed strong antibacterial activities against the tested bacteria with MIC values ranging from 0.5 to 6.3 μ M. Compounds **15** and **20** exhibited strong antibacterial activities against *B. subtilis*, *S. aureus*, and *K. pneumonia* with MIC values ranging from 0.5 to 6.3 μ M, whereas the assay revealed antibacterial activities of **17** and **19** against the tested positive bacteria with MIC values ranging from 0.5 to 25 μ M. However, **18**, **21**, and **22–24a/b** didn't exhibit any antibacterial activities. Notably, the antibacterial activity was investigated for the first time among the isolated PBDEs with the exception of **14**, **18**, and **21**.

Table 3.3 Antibacterial activities of isolated compounds from *Arenosclera* sp.

Samples	Minimum Inhibitory Concentration (MIC) (μ M)			
	<i>Bacillus subtilis</i>	<i>Staphylococcus aureus</i>	<i>Klebsiella pneumoniae</i>	<i>Escherichia coli</i>
13	3.1	6.3	6.3	>50
14	0.5	0.5	3.1	6.3
15	0.8	0.5	6.3	12.5
16	12.5	25	25	>50
17	0.5	1.6	>50	>50
18	>50	>50	>50	>50
19	0.5	0.8	>50	>50
20	3.1	6.3	6.3	>50
21	>50	>50	>50	>50
22	>50	>50	>50	>50
23	>50	>50	>50	>50
24a/b	>50	>50	>50	>50
Ampicillin ^a	3.1	3.1	—	—
Kanamycin ^b	—	—	3.1	6.3

^a positive control for Gram-positive bacteria. ^b positive control for Gram-negative bacteria.

3.7 Summary of chapter 3

One new polybrominated diphenyl ether, tribromiododiphenyl ether (**13**), together with eight known PBDEs (**14–21**) and three known sterols (**22–24-a/b**), were isolated from the MeOH extract of the Vietnamese marine sponge *Arenosclera* sp..

The secondary metabolites with iodine are rarely seen even in the metabolites isolated from the marine natural resources.

The new compound **13** only showed moderate cytotoxic activity against MCF-7 with the IC₅₀ value of 35.6 μ M. However, **13** possessed strong antibacterial activities against *B. subtilis*, *S. aureus*, and *K. pneumoniae*, with the MIC values ranging between 3.1–6.3 μ M. Most of the known PBDEs, especially **14**, displayed strong anti-bacterial activities against the tested four bacteria with the MIC values ranging from 0.5–6.3 μ M, as well as strong cytotoxic activities against the tested three cancer cell lines with the IC₅₀ values ranging from 5.0–7.5 μ M.

In this study, these occurrences in *Arenosclera* sp. provide insight into not only the bioactive compounds in the Vietnamese *Arenosclera* sp. marine sponge, but also the biodiversity of *Arenosclera* species from the view point of their chemical constituents.

Chapter 4 Constituents of *Gelliodes* sp. collected in Vietnam and their cytotoxic activities

4.1 Introduction

The genus *Gelliodes* consists of sponges with elastic texture and belongs to the Niphatidae family. The *Gelliodes* sponges appear as dark purple, grey, dark brownish colors and have straight or slightly curved spicules about $160 \pm 180 \times 5 \mu\text{m}$ ⁸¹. The sponges from this genus are distributed in several regions around the Indian Ocean, West Pacific Ocean, New Guinea, New Caledonia, and South China Sea⁵².

To the best of our knowledge, only twelve carotenoids including a new aldehydic carotenoid, gelliodesxanthin, were isolated from *G. callista* as the chemical components⁸². On the other hand, there are reports on the bioactive chemical components of streptomycetes and fungus isolated from marine sponge *Gelliodes* sp.. Investigation of the chemical components of the *Streptomyces* sp. LS298 from *G. carnosa* resulted in the isolation of two new compounds, a new echinomycin derivative and a new cyclic dipeptide derivative along with three known compounds, tirandamycines A and B and staurosporine⁸³. In addition, the chemical studies of two fungi, *Nectria* sp. HLS206 and *Hypocreales* sp., associated with *G. carnosa* have led to isolation of a set of three α -pyrone derivatives⁸⁴ and a set of three cadinane-type sesquiterpenes, ergosterol, two α -pyrone derivatives, and coumaric acid, respectively⁸⁵. These results indirectly suggested that Vietnam marine sponge *Gelliodes* sp. contains these type compounds in nature. In the course of our screening, the MeOH extract of *Gelliodes* sp. collected from Vietnam showed cytotoxic activities against HeLa, MCF-7, and A549. Thus, further study was performed to investigate the chemical constituents of *Gelliodes* sp. with cytotoxic compounds.

4.2 Extraction and isolation

Gelliodes sp. (611 g dry weight) was macerated in MeOH (2 L $\times$ 3) for 2 h with sonication. This extract (62.3 g) was evaporated under vacuum and partitioned with *n*-hexane to obtain the *n*-hexane fraction. The defatted extract was subjected to HP-20 column chromatography using MeOH and H₂O to give H₂O (containing mainly salts, not evaporated), 50% MeOH (1.1 g), and MeOH (2.5 g) fractions, respectively. The MeOH fraction was subjected to normal phase silica gel open column chromatography using solvent systems composed of CHCl₃/MeOH with increasing polarities. In total, 40 fractions of 300 mL each were collected and pooled into 10 combined fractions after TLC profiling (Figure 4.2). These fractions were further subjected to series of chromatographic separation, which led to isolation of eleven compounds including one new decenoic acid (**25**), a natural new indole alkaloid, 3-(propan-2-ylidene) indolin-2-one (**26**)⁸⁶, three known furanoterpenoids, sarcotin H (**27**)⁸⁷, (7*E*, 12*E*, 20*Z*, 18*S*)-variabilin (**28**)⁸⁸, and sulawesin C (**29**)⁸⁹, and two known indole alkaloids, 1*H*-indole-3-carboxylic acid (**30**)⁹⁰ and 3-(hydroxyacetyl)indole (**31**)⁹¹ (Figure 4.3).

Figure 4.1 *Gelliodes* sp.

Figure 4.2 Extraction and isolation of compounds from MeOH extract of *Gelliodes* sp.

Figure 4.3 Structures of isolated compounds **25–31** from *Gelliodes* sp.

4.3 Structure elucidation of new compounds

4.3.1 Gelliodesinic acid (**25**)

Compound **25** obtained as colorless oil. The HR-ESI-MS displayed a pseudo-molecular ion peak at m/z 221.1151 $[M+Na]^+$ (calcd. for $C_{11}H_{18}O_3Na$, 221.1154) and the molecular formula of **25** was established as $C_{11}H_{18}O_3$, indicating three degrees of unsaturation. The UV spectrum showed maximum absorptions at 206, 212, 218, 228 and 294 nm. The IR spectrum exhibited absorption bands at 3419 cm^{-1} (O–H), 2925 cm^{-1} , 2854 cm^{-1} (C–H), 1756 cm^{-1} (C=O), 1712 cm^{-1} (COOH), and 1654 cm^{-1} (C=C).

The 1H NMR spectrum of **25** showed signals of an olefinic proton at δ_H 5.70 (s, H-2), two methyl protons at δ_H 1.92 (s, H-11) and 1.05 (t, $J = 7.4$ Hz, H-10), and five methylene protons at δ_H 2.64 (t, $J = 7.8$ Hz, H-4), 2.45 (t, $J = 7.0$ Hz, H-7), 2.42 (t, $J = 7.4$ Hz, H-9), 1.63 (m, H-6), and 1.48 (m, H-5). The ^{13}C NMR and HMQC spectra of **25** revealed the presence of two carbonyl carbons at δ_C 211.8 (C-8) and 169.4 (C-1), two olefinic carbons at δ_C 163.1 (C-3) and 115.4 (C-2), five methylene carbons at δ_C 41.9 (C-7), 35.9 (C-9), 33.0 (C-4), 27.6 (C-5), and 23.6 (C-6), and two methyl carbons at δ_C 25.3 (C-11) and 7.7 (C-10) (Table 4.1, Figure 4.4, and 4.5) in **25**. These NMR data were similar to those of synthetic compound, (Z)-8-hydroxy-3-methyldec-2-enoic acid, reported by Emura *et al*⁹².

The 1H – 1H COSY correlations $[C(4)H_2-C(5)H_2-C(6)H_2-C(7)H_2]$ and HMBC correlation from H₂-6 to C-4 and C-8 indicated the presence of a linear chain moiety (C-4–C-7) in **25**. The spin systems $[C(9)H_2-C(10)H_3]$ in the 1H – 1H COSY spectrum and the cross peaks from H₃-10, H₂-9, and H₂-7 to C-8 in the HMBC spectrum revealed the presence of a propan-1-one moiety (C-8–C-10). The HMBC correlations from H₂-4 to C-2 and C-3, from H₃-11 to C-2 and C-3, from H₂-4 to C-11 and C-4, and from H-2 to C-4

and C-11 indicated the location of the olefinic bond at C-2 and C-3. Following the molecular formula of **25**, remaining carboxyl group should be connected to C-2 (Figure 4.6). The geometry of the trisubstituted olefin (C-2–C-3) was assigned as *Z*-configuration by the NOESY correlation of H-2/H₃-11 (Figure 4.6). Compound **25** was determined to be decenoic acid derivative and named gelliodesinic acid. Compound **25** was suggested as a biosynthetic precursor of (2*Z*,8*R*)-8-hydroxy-3-methyl-2 -decenoic acid isolated from *ircinia echinata*⁹³.

Figure 4.4 ^1H NMR (500 MHz) spectrum of gelliodesinic acid (**25**)

Figure 4.5 ^{13}C NMR (125 MHz) spectrum of gelliodesinic acid (**25**)

Figure 4.6 HMBC (arrows), ^1H - ^1H COSY (bold lines), and NOESY (dotted arrow) of gelliodesinic acid (**25**)

Table 4.1 ^1H (500 MHz) and ^{13}C NMR (125 MHz) data of **25** in CDCl_3 (δ in ppm and J values in (Hz) in parentheses)

Position	25	
	δ_{H}	δ_{C}
1		169.4 *
2	5.70 (s)	115.4
3		163.1
4	2.64 (t, 7.8)	33.0
5	1.48 (m)	27.6
6	1.63 (m)	23.6
7	2.45 (t, 7.0)	42.0
8		211.8
9	2.42 (t, 7.4)	35.9
10	1.05 (t, 7.4)	7.7 *
11	1.92 (s)	25.3

* ^{13}C chemical shifts determined from HMQC and HMBC.

4.4 Cytotoxic activities of isolated compounds from *Gelliodes* sp.

All isolated compounds (**25–31**) were tested for cytotoxic activities against human cervical cancer cell line HeLa, human breast cancer cell line MCF-7, and human lung cancer cell line A549 at the concentrations of 10, 50, and 100 μM using the MTT assay.

The isolated furanoterpenoids **27–29** showed cytotoxic activities against three cancer cell lines. Compound **25** didn't show cytotoxic activity against three tested cell lines. Compounds **27–29** displayed moderate cytotoxic activities against the three tested cancer cell lines with the IC_{50} values ranging from 13.2 to 75.5 μM . However, the isolated indole alkaloids **26**, **30**, and **31** didn't show any cytotoxic activities except for **30**. The known compound **30** displayed weak cytotoxic activity against A549 with the IC_{50} value of 89.2 μM .

Table 4.2 Cytotoxic activities of isolated compounds from *Gelliodes* sp.

Samples	IC ₅₀ (μM)		
	HeLa	MCF-7	A549
25	> 100	> 100	> 100
26	> 100	> 100	> 100
27	41.8	29.3	23.6
28	35.4	25.0	75.5
29	69.0	54.2	58.6
30	> 100	> 100	89.2
31	> 100	> 100	> 100
5-FU ^a	43.1	46.2	47.5

^a 5-Fluorouracil: positive control.

4.5 Summary of chapter 4.

One new fatty acid derivative, gelliodesinic acid (**25**), together with three known furanoterpenoids (**27–29**) and three known indole derivatives (**26**, **30**, and **31**) were isolated from the MeOH extract of the Vietnamese marine sponge *Gelliodes* sp.. Compound **26** was reported as natural compound for the first time.

Analyses of the isolated furanoterpenoids (**27–29**) suggested that they possessed moderate cytotoxic activities against tested three cancer cell lines. From these results, the furanoterpenoids could be potential candidates for cytotoxic activities.

The results provided new insights into not only chemical components of Vietnamese *Gelliodes* sp., but also chemodiversity of furanoterpenoids and indole derivatives in marine sponges.

■ Discussion

In the search of bioactive compounds from marine sponges, I have isolated several types of compounds. Among them, the isolation of polyacetylene amide and PBDEs from marine sponges *S. siphonella* and *Arenosclera* sp., respectively, were somehow suggested to be due to the presence of cyanobacteria in several reported literatures previously. Notable, *N*-acylated tryptamides or phenethylamides were reported to be isolated from Cyanobacterium *Lyngbya majuscula* collected from tropical and sub-tropical shallow water, while the discovery of PDBEs compounds were found within the cyanobacteria, *Hormscilla spngeliae* (formerly *Oscillatoria spongelliae*), *Moorea producens*, *Pleurocapsa* sp., and *Rivularia* sp. that were found in marine sponges ^{94, 95}. These interesting observations may suggest that the discovery of similar types of compounds in my study might also be due to the presence of cyanobacterium in the marine sponges. However, different species of cyanobacterium might produce different classes of secondary metabolites due to a variety of environmental factors as the biosynthetic origin is still remained unclear at this moment.

In terms of bioactivity, polyacetylene compounds obtained from *Panax ginseng* were previously reported to possess cytotoxic activities against normal cell line, Mesothelial cells ⁹⁶. PBDEs compounds were also reported in the previous literature to have cytotoxic activities against Human L02 cell line ⁹⁷. This may suggest that the similar types of compounds discovered in my study might also possess cytotoxicity against normal cell lines.

■ Conclusion

In this study, the chemical constituents of a marine sponge *Siphonochalina siphonellla* collected from Egypt and two marine sponges *Arenosclera* sp. and *Gelliodes* sp. collected from Vietnam were investigated. Total 31 compounds (**1–31**) including ten new compounds (**1–8**, **13**, and **25**) named siphonellamides A–E (**1–5**), siphonellanol A–C (**6–8**), tribromiododiphenyl ether (**13**), and gelliodesinic acid (**25**) were isolated. The indole acetylene amide types of compounds **2–4** were isolated from natural resource for the first time, while PBDEs (**13–21**) were isolated for the first time from *Arenosclera* sp.. On the other hand, chemical constituents of Vietnamese *Arenosclera* sp. and *Gelliodes* sp. have not been thoroughly investigated.

The isolated compounds from this study were classified into five types on the skeletons, polyacetylenes (**1–4**, **6–8**, and **10–12**), PBDEs (**13–21**), furanoterpenoids (**27–29**), indole alkaloids (**26**, **30**, and **31**), and lipids (**5**, **9**, and **22–25**). Among these skeletons, polyacetylene, PBDEs, and furanoterpenoids showed cytotoxic activities. These results suggested that these three skeletons could be attractive candidates for further investigation as anti-cancer drugs.

The present study in conjugated with the cytotoxic activities against human cancer cell lines provided new insight into not only chemical components of the Egyptian and Vietnamese marine sponges, but also chemodiversity of the isolated compounds in marine sponges.

■ Experimental

I. General Experimental Procedures

Infrared spectra were recorded with KBr pellets on a JASCO FT/IR-460 Plus spectrometer (Japan). UV-spectra were recorded on a NANODROP 2000C spectrophotometer (Thermo Fisher Scientific, USA). ^1H and ^{13}C NMR spectra were recorded on JEOL ECA500II or JEOL ECA400II spectrometers (Japan). COSY and TOCSY NMR spectra were measured on a Bruker Avance III 800 spectrometer. Chemical shift values are expressed in δ (ppm) downfield from TMS, as an internal standard. HR-ESI-MS, LR-ESI-MS and ESI-MS/MS data were obtained in positive and negative mode on SHIMADZU LC-MS-IT-TOF (Japan) and Agilent Technologies 6420 Triple Quad LC/MS spectrometers (ESI voltage: + 3.0 kV), respectively. Open column chromatography was performed with normal phase silica gel (silica gel 60 N, spherical, neutral, 40–50 μM) (Kanto Chemicals, Japan) and reversed phase silica gel (Cosmosil 75C₁₈-OPN) (Nacalai Tesque, Japan). TLC was performed on silica gel GF₂₅₄ pre-coated plates (Merck, USA), with detection by visualization with a UV lamp at 254 and 365 nm, followed by spraying with a *p*-anisaldehyde stain solution (Nacalai Tesque, Kyoto, Japan) and heating to 120°C. HPLC was performed with an Agilent Technologies 1260 quaternary pump with an Agilent Infinity II 1260 UV/Visible detector and silica gel 60 column (21.5 mm I.D. $\times$ 300 mm) (TOSOH TSK, Japan) was used for the normal phase HPLC column chromatography and an Agilent Technologies 1260 quaternary pump with a HEWLETT PACKARD 1100 detector and Agilent Infinity II 1260 UV/Visible detector and an OSAKA SODA C₁₈ CAPCELL PAK (10 mm I.D $\times$ 250 mm) and 5C₁₈-AR- II (10 mm I.D $\times$ 250 mm) (Nacalai Tesque, Japan) was used for the reversed-phase HPLC column chromatography.

Cytotoxic activity of isolated compounds

The isolated compounds were evaluated cytotoxic activities using the 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Nacalai Tesque, Japan) assay, according to the published procedure. α -Minimum essential medium with l-glutamine and phenol red (α -MEM; Wako, Japan) were used to culture the A549, HeLa, and MCF-7 cancer cell lines. All media were supplemented with 10% fetal bovine serum (FBS; Sigma, USA) and 1% antibiotic antimycotic solution (Sigma, USA). For the MCF-7 cells, the growth medium was supplemented with 1% 0.1 mM non-essential amino acids (NEAA; Gibco, USA) and 1% 1 mM sodium pyruvate (Gibco, USA). Each cancer cell line was seeded in 96-well plates (1×10^4 cells per well) and incubated in the respective medium at 37°C, under 5% CO₂ and 95% air atmosphere, for 24 h. After the cells were washed with phosphate-buffered saline (PBS), different concentrations of the tested samples (10, 50, and 100 μ M) were added. After 72 h incubation, the cells were washed with PBS and 100 μ L of medium containing 10 μ L of MTT solution (5 mg/mL) was added to each well and incubated for 3 h. Subsequently, the absorbance of each well was measured at a 570 nm wavelength. 5-Fluorouracil (Wako, Japan) was used as a positive control.

2 Experimental detail of chapter 2

Marine sponge material

S. siphonella was collected in January 2018 from the reefs located southwest of Magawish Island, Hurghada, Egypt, at a depth of 7 m and immediately soaked in MeOH. A vouchered specimen was deposited in the Red Sea Marine Parks, Hurghada, Egypt.

Extraction and isolation

S. siphonella (600 g wet weight) was macerated in MeOH (2 L $\times$ 3) for 2 h with sonication. The extract was evaporated under vacuum and the remaining aqueous residue was successively partitioned with CHCl₃, EtOAc, and BuOH. The CHCl₃ fraction (4.5 g) was subjected to normal phase silica gel open column chromatography, using solvent systems composed of *n*-hexane/EtOAc and EtOAc/MeOH, with increasing polarities. In total, 190 fractions of 30 mL each were collected and pooled into 21 combined fractions after TLC profiling.

Compounds **1** (4.3 mg) and **10** (1.2 mg) were acquired from the combined fraction 17 (83.2 mg) which obtained *n*-hexane/EtOAc (3:7) using ODS column chromatography eluted with MeCN/H₂O (3:1). Compounds **2** (1.5 mg, *t_R* 90 min), **3** (0.6 mg, *t_R* 83 min), and **4** (0.7 mg, *t_R* 93 min) were isolated from fraction 19 (105.1 mg) using reverse-phase semi-preparative HPLC equipped OSAKA SODA C₁₈ CAPCELL PAK (10 mm I.D $\times$ 250 mm) column with gradient elution by MeCN/H₂O (3:2, 1–60 min), MeCN/H₂O (7:3, 60–120 min), and MeCN (120–150 min) at flow rate 2 mL/min, monitored at 254 nm. Compounds **5** (4.8 mg) and **9** (1.2 mg) were purified from combined fraction 18 (35.4 mg) using ODS column chromatography eluted with MeCN/H₂O (3:1).

Fractions 15 and 16 (239.9 mg) were combined and then performed silica gel column chromatography using *n*-hexane/EtOAc (8.5:1.5) to yield the F_{1-5} sub fractions. Compounds **6** (3.4 mg, t_R 72 min), **7** (1.2 mg, t_R 91 min), **8** (1.6 mg, t_R 108 min), and **12** (4.5 mg, t_R 60 min) were purified from F_3 sub fraction (70 mg) using normal phase preparative HPLC equipped silica 60 column (21.5 mm I.D. $\times$ 300 mm) with elution by *n*-hexane/EtOAc (4:1) isocratic solvent system (flow rate 6 mL/min; running time 120 min) and monitored at a 254 nm wavelength. Compound **11** (100 mg) was isolated from combined fraction 14 (316.2 mg) which obtained from *n*-hexane/EtOAc (1:1) using silica gel column chromatography eluted with *n*-hexane/EtOAc (8.5:1.5).

Ozonolysis

A solution of **5** (0.2 mg) in dichloromethane (1.0 mL) was treated with O_3 at $-78^\circ C$ for 15 min, and then triphenylphosphine (0.5 mg) was added to the reaction mixture at $-78^\circ C$ for 3 h. The product was warmed to room temperature and evaporated in vacuo, and the resultant solution was subjected to ESI-MS and ESI-MS/MS (CID 30 eV) in the positive mode without further purification.

Siphonellamide A (1): yellowish oil. IR (KBr) ν_{max} : 3434, 3287, 2926, 2854, 2352, 1648, 1539, 1457, 755, 664 cm^{-1} . UV (MeOH) λ_{max} (log ϵ): 211 (4.0), 215 (4.0), 218 (4.0), 228 (3.9). 1H NMR (400 MHz, $CDCl_3$) and ^{13}C NMR (100 MHz, $CDCl_3$): see Table 2.1. HR-ESI-MS (+): m/z 414.2762 $[M+Na]^+$ (calcd. for $C_{27}H_{37}NONa$, 414.2773). ESI-MS/MS (daughter ion, CID 30 eV): m/z 414 ($[M+Na]^+$, 100), 200 ($[M-C_{14}H_{23}]^+$, 20), 121 ($[M+H-C_{19}H_{27}O]^+$, 18), 105 ($[M-C_{19}H_{28}NO]^+$, 18).

Siphonellamide B (2): yellowish oil. IR (KBr) $\nu_{\max}$: 3422, 3276, 2926, 2860, 2356, 1728, 1642, 1548, 1457, 1227, 1109, 752 cm^{-1} . UV (MeOH) $\lambda_{\max}$ (log ϵ): 218 (4.0), 227 (4.0), 290 (3.3). ^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3): see Table 2.2. HR-ESI-MS (+): m/z 453.2865 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{29}\text{H}_{38}\text{N}_2\text{ONa}$, 453.2882). ESI-MS/MS (daughter ion, CID 15 eV): m/z 431 ($[\text{M}+\text{Na}]^+$, 100), 187 ($[\text{M}-\text{C}_{18}\text{H}_{27}]^+$, 10), 160 ($[\text{M}+\text{H}-\text{C}_{19}\text{H}_{27}\text{O}]^+$, 11), 144 ($[\text{M}-\text{C}_{19}\text{H}_{30}\text{NO}]^+$, 82).

Siphonellamide C (3): yellowish oil. IR (KBr) $\nu_{\max}$: 3404, 3299, 2926, 2860, 2357, 1734, 1646, 1548, 1454, 1229, 1109, 752 cm^{-1} . UV (MeOH) $\lambda_{\max}$ (log ϵ): 219 (4.0), 227 (4.0), 290 (3.3). ^1H NMR (400 MHz, CDCl_3): see Table 2.3. ^{13}C NMR was deduced from HMQC and HMBC: see Table 2.3. HR-ESI-MS (+): m/z 451.2712 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{29}\text{H}_{36}\text{N}_2\text{ONa}$, 451.2725). ESI-MS/MS (daughter ion, CID 45 eV): m/z 429 ($[\text{M}+\text{H}]^+$, 20), 144 ($[\text{M}-\text{C}_{19}\text{H}_{26}\text{NO}]^+$, 100), 130 ($[\text{M}-\text{C}_{20}\text{H}_{28}\text{NO}]^+$, 15), 116 ($[\text{M}-\text{C}_{21}\text{H}_{30}\text{NO}]^+$, 10), 105 ($[\text{M}-\text{C}_{21}\text{H}_{27}\text{N}_2\text{O}]^+$, 15).

Siphonellamide D (4): yellowish oil. IR (KBr) $\nu_{\max}$: 3414, 3299, 2927, 2865, 1731, 1645, 1545, 1448, 1232, 1106, 752 cm^{-1} . UV (MeOH) $\lambda_{\max}$ (log ϵ): 220 (4.0), 228 (4.0), 290 (3.2). ^1H NMR (400 MHz, CDCl_3): see Table 2.4. ^{13}C NMR was deduced from HMQC and HMBC: see Table 2.4. HR-ESI-MS (+): m/z 441.2881 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{28}\text{H}_{38}\text{N}_2\text{ONa}$, 441.2882). ESI-MS/MS (daughter ion, CID 35 eV): m/z 441 ($[\text{M}+\text{Na}]^+$, 100), 163 ($[\text{M}-\text{C}_{16}\text{H}_{19}\text{N}_2\text{O}]^+$, 10), 144 ($[\text{M}-\text{C}_{18}\text{H}_{28}\text{NO}]^+$, 38).

Siphonellamide E (5): yellowish oil. IR (KBr) $\nu_{\max}$: 3406, 2925, 1642, 1542,

1457, 1094, 744 cm^{-1} . UV (MeOH) λ_{max} ($\log \epsilon$): 215 (4.1), 219 (4.2), 226 (4.2), 282 (3.6). ^1H NMR (400 MHz) and ^{13}C NMR (100 MHz, CDCl_3): see Table 2.5. HR-ESI-MS (+): m/z 419.3025 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{26}\text{H}_{40}\text{N}_2\text{ONa}$, 419.3038). ESI-MS/MS (daughter ion, CID 40 eV): m/z 419 ($[\text{M}+\text{Na}]^+$, 60), 159 ($[\text{M}-\text{C}_{16}\text{H}_{29}\text{O}]^+$, 60), 144 ($[\text{M}-\text{C}_{16}\text{H}_{30}\text{NO}]^+$, 100).

Siphonellanol A (6): yellowish oil; IR (KBr) ν_{max} : 3432, 3303, 2928, 2865, 2355, 2213, 1717, 1639, 1533, 1554, 1018, 759 cm^{-1} ; UV (MeOH) λ_{max} ($\log \epsilon$): 213 (4.0), 218 (4.0), 229 (4.0). ^1H NMR (500 MHz, CDCl_3) and ^{13}C NMR (125 MHz, CDCl_3): see Table 2.6. HR-ESI-MS (+): m/z 325.1543 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{22}\text{H}_{22}\text{ONa}$, 325.1568). ESI-MS/MS (daughter ion, CID 32 eV): m/z 325 ($[\text{M}+\text{Na}]^+$, 100), 237 ($[\text{M}-\text{C}_5\text{H}_5]^+$, 35), 208 ($[\text{M}+\text{H}-\text{C}_6\text{H}_7\text{O}]^+$, 35), 180 ($[\text{M}+\text{H}-\text{C}_8\text{H}_{11}\text{O}]^+$, 15).

Siphonellanol B (7): yellowish oil; IR (KBr) ν_{max} : 3413, 3293, 2931, 2854, 2365, 2213, 1720, 1649, 1528, 1454, 1218, 1018, 759 cm^{-1} ; UV (MeOH) λ_{max} ($\log \epsilon$): 214 (4.0), 219 (4.0), 228 (4.0). ^1H NMR (500 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3): see Table 2.7. HR-ESI-MS (+): m/z 329.1858 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{22}\text{H}_{26}\text{ONa}$, 329.1881). ESI-MS/MS (daughter ion, CID 20 eV): m/z 329 ($[\text{M}+\text{Na}]^+$, 100), 261 ($[\text{M}-\text{C}_2\text{H}_5\text{O}]^+$, 40), 180 ($[\text{M}+\text{H}-\text{C}_8\text{H}_{15}\text{O}]^+$, 35).

Siphonellanol C (8): yellowish oil; IR (KBr) ν_{max} : 3431, 3303, 2927, 2865, 2349, 2213, 1709, 1640, 1533, 1445, 1018, 758 cm^{-1} ; UV (MeOH) λ_{max} ($\log \epsilon$): 213 (3.9), 219 (3.9), 229 (4.0). ^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3): see Table 2.8. HR-ESI-MS (+): m/z 327.1708 $[\text{M}+\text{Na}]^+$ (calcd. for $\text{C}_{22}\text{H}_{24}\text{ONa}$, 327.1725). ESI-MS/MS (daughter ion, CID 19 eV): m/z 327 ($[\text{M}+\text{Na}]^+$, 100), 197 ($[\text{M}-\text{C}_8\text{H}_{11}]^+$, 35), 180

([M+H-C₈H₁₃O]⁺, 40).

***N*-[2''-(1'*H*-Indol-3'-yl)ethyl]-hexadecanamide (9)**: yellowish oil; ¹H NMR (500 MHz, DMSO-*d*₆) δ: 2.03 (t, 2H, *J* = 7.7 Hz, H-2), 1.46 (m, 2H, H-3), 1.23 (overlapped, br s, 24H, H-4–H-15), 0.85 (t, *J* = 6.8 Hz, 3H, H-16), 10.8 (br s, 1H, 1-NH), 7.11 (d, 1H, *J* = 2.6 Hz, H-2'), 7.53 (d, 1H, *J* = 7.9 Hz, H-4'), 6.96 (m, 1H, H-5'), 7.06 (m, 1H, H-6'), 7.32 (d, 1H, *J* = 8.3 Hz, H-7'), 3.31 (m, 2H, H-1''), 2.79 (t, 2H, *J* = 7.9 Hz, H-2''), 7.87 (m, 1H, -NH).

Callyspongamide A (10): yellowish oil; ¹H NMR (500 MHz, CDCl₃) δ: 2.31 (t, *J* = 7.1 Hz, 2H, H-2), 2.46 (tt, *J* = 7.0, 2.5 Hz, 1H, H-3), 2.08 (tt, *J* = 7.2, 2.4 Hz, 1H, H-6), 1.44 (m, 2H, H-7), 1.30 (overlapped, m, 8H, H-8–H-11), 1.41 (m, 2H, H-12), 2.34 (m, 2H, H-13), 5.99 (td, *J* = 10.8, 7.5 Hz, 1H, H-14), 5.44 (br d, *J* = 10.8 Hz, 1H, H-15), 3.07 (d, *J* = 2.2 Hz, 1H, H-17), 5.72 (br s, 1H, -NH), 3.53 (q, *J* = 7.0 Hz, 2H, H-1'), 2.82 (t, *J* = 7.0 Hz, 2H, H-2'), 7.21 (overlapped, d, *J* = 7.6 Hz, 2H, H-4' and H-8'), 7.32 (overlapped, t, *J* = 7.6 Hz, 2H, H-5' and H-7'), 7.24 (t, *J* = 7.4 Hz, 1H, H-6').

Dehydroisophonochalynol (11): yellowish oil; ¹H NMR (400 MHz, CDCl₃) and ¹³C NMR (100 MHz, CDCl₃): see Table 2.6.

Siphonochalynol (12): yellowish oil; ¹H NMR (500 MHz, CDCl₃) δ: 4.25 (t, *J* = 2.2 Hz, 2H, H-1), 2.30 (q, *J* = 7.1 Hz, 2H, H-4), 1.52 (overlapped, m, 10H, H-5, H-6, H-7, H-16, and H-17), 2.35 (overlapped, q, *J* = 7.2 Hz, 4H, H-8 and H-15), 2.23 (m, 2H, H-18), 5.97 (td, *J* = 10.9, 7.5 Hz, 1H, H-19), 5.47 (d, *J* = 10.9 Hz, 1H, H-20), 3.08 (d, *J*

= 2.3 Hz, 1H, H-22).

3 Experimental detail of chapter 3

Marine sponge material

The *Arenosclera* sp. marine sponges were collected from Hon Thom, Phu Quoc, Vietnam, in June 2017, and positively identified by Quang Minh Thai at the Vietnam Academy of Science and Technology, Institute of Oceanography, Nha Trang (Vietnam). A voucher specimen (E55481) was deposited at the Oceanography Museum, Institute of Oceanography in Nha Trang, Vietnam.

Extraction and isolation

Arenosclera sp. (474 g dry weight) was macerated in MeOH (2 L x 3) for 2 h with sonication. The crude extract (79.7 g) was evaporated under vacuum. The crude extract was subjected to HP-20 column chromatography using MeOH and H₂O to give H₂O, 50% MeOH (0.6 g), and MeOH (52.5 g) fractions, respectively. MeOH fraction was subjected to normal phase silica gel open column chromatography using solvent systems composed of *n*-hexane/EtOAc and EtOAc/MeOH with increasing polarities. In total, 83 fractions of 500 mL each was collected and pooled into 15 combined fractions after TLC profiling.

Compounds **13** (1.4 mg), **16** (10.7 mg), **17** (5.8 mg), **19** (13.2 mg), and **20** (7.7 mg) were acquired from combined fraction 2 (224.1 mg) using *n*-hexane/EtOAc (50:1–1:1) with increasing polarities and preparative reverse phase TLC with acetone/MeOH/MeCN/water (6:1:1:1) solvent system. Compounds **14** (8.7 mg) and **18** (2.4 mg) were isolated from combined fraction 3 (15.9 g) which obtained from *n*-hexane/EtOAc (9:1) using Sephadex LH-20 column chromatography eluted with acetone.

Compound **15** (5.8 mg) was acquired from combined fraction 4 (7.1 g) which obtained from *n*-hexane/EtOAc (7:1) using Sephadex LH-20 column chromatography eluted with MeOH. Compound **21** was isolated from combined fraction 1 (142.7 mg) which obtained from *n*-hexane/EtOAc (50:1) using silica gel column chromatography with gradient eluted with *n*-hexane/EtOAc (20:1–1:1). The combined fractions 6 (1.16 g) and 9 (337.5 mg) were separately subjected to silica gel column chromatography with gradient elution by *n*-hexane/EtOAc (50:1–1:1), and preparative normal phase TLC to yield **22** (5.4 mg), **23** (3.7 mg), and **24a/b** (1.7 mg).

Tribromoiiododiphenyl ether (13): white powder. IR (KBr) ν_{max} : 3428, 2920, 1577, 1465, 1093, 690, and 550 cm^{-1} . UV (MeOH) λ_{max} (log ϵ): 229 (4.1), 283 (3.4), and 290 (3.4). ^1H NMR (500 MHz, CDCl_3) and ^{13}C NMR (125 MHz, CDCl_3): see Table 3.1. HR-ESI-MS (-): 550.6822 $[\text{M}+6\text{-H}]^-$ (30), 548.6853 $[\text{M}+4\text{-H}]^-$ (83), 546.6889 $[\text{M}+2\text{-H}]^-$ (74), 544.6859 $[\text{M-H}]^-$ (26) (calcd. for $\text{C}_{12}\text{H}_5\text{O}_2\text{Br}_3\text{I}$, 544.6890), 520.9106 (83), 498.9194 (43), 384.9346 (95), 362.9321 (13), 339.2044 (30), 325.1827 (56), 311.1689 $[\text{M-H-C}_6\text{H}_3\text{OBr}_2]^-$ (48), 293.01774 (13), 265.1484 (22), and 248.9590 $[\text{M-H-C}_6\text{H}_3\text{OBrI}]^-$ (100).

2-(2',4'-Dibromophenoxy)-4,6-dibromophenol (14): white powder. ^1H NMR (500 MHz, CDCl_3) δ : 7.22 (d, $J = 2.3$ Hz, 1H, H-3), 7.35 (d, $J = 2.3$ Hz, 1H, H-5), 5.54 (br s, 1H, 1-OH), 7.77 (d, $J = 2.3$ Hz, 1H, H-3'), 7.30 (dd, $J = 9.2, 2.3$ Hz, 1H, H-5'), 6.45 (d, $J = 9.2$ Hz, 1H, H-6'). ^{13}C NMR (125 MHz, CDCl_3) δ : 138.6 (C-1), 150.3 (C-2), 119.6 (C-3), 117.2 (C-4), 127.7 (C-5), 119.8 (C-6), 152.0 (C-1'), 112.7 (C-2'), 136.2 (C-3'), 116.0 (C-4'), 131.5 (C-5'), 115.9 (C-6').

2-(4'-Bromophenoxy)-3,5-dibromophenol (15): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 7.34 (d, $J = 2.3$ Hz, 1H, H-4), 7.20 (d, $J = 2.3$ Hz, 1H, H-6), 5.51 (br s, 1H, 1-OH), 6.78 (dd, $J = 9.2, 2.3$ Hz, 2H, H-2' and H-2''), 7.42 (dd, $J = 9.2, 2.3$ Hz, 2H, H-3' and H-3''). ^{13}C NMR (125 MHz, CDCl_3) δ : 150.5 (C-1), 138.3 (C-2), 117.5 (C-3), 127.7 (C-4), 119.4 (C-5), 119.3 (C-6), 155.3 (C-1'), 116.9 (C-2' and C-2''), 132.9 (C-3' and C-3''), 115.8 (C-4').

5-Bromo-2-(2',4'-dibromophenoxy) phenol (16): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 7.31 (dd, $J = 8.6, 1.7$ Hz, 1H, H-3), 6.76 (d, $J = 8.6$ Hz, 1H, H-4), 6.77 (s, 1H, H-6), 5.90 (br s, 1H, 1-OH), 7.79 (d, $J = 2.3$ Hz, 1H, H-3'), 7.39 (dd, $J = 8.6, 2.3$ Hz, 1H, H-5'), 6.83 (d, $J = 8.6$ Hz, 1H, H-6'). ^{13}C NMR (125 MHz, CDCl_3) δ : 144.4 (C-1), 143.7 (C-2), 128.3 (C-3), 117.6 (C-4), 110.4 (C-5), 121.2 (C-6), 152.4 (C-1'), 115.1 (C-2'), 136.2 (C-3'), 117.1 (C-4'), 131.7 (C-5'), 120.6 (C-6').

2-(2',4'-Dibromophenoxy)-5,6-dibromo-3-chlorophenol (17): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 7.55 (s, 1H, H-4), 6.06 (br s, 1H, 1-OH), 7.78 (d, $J = 2.3$ Hz, 1H, H-3'), 7.29 (dd, $J = 9.2, 2.3$ Hz, 1H, H-5'), 6.43 (d, $J = 9.2$ Hz, 1H, H-6'). ^{13}C NMR (125 MHz, CDCl_3) δ : 147.1 (C-1), 139.1 (C-2), 122.7 (C-3), 127.7 (C-4), 119.9 (C-5), 115.9 (C-6), 152.2 (C-1'), 112.7 (C-2'), 136.1 (C-3'), 115.9 (C-4'), 131.3 (C-5'), 115.8 (C-6').

2-(2',4'-Dibromophenoxy)-3,5-dibromophenol (18): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 6.81 (d, $J = 2.3$ Hz, 1H, H-4), 7.44 (d, $J = 2.3$ Hz, 1H, H-6), 5.94 (br

s, 1H, 1-OH), 7.82 (d, $J = 2.3$ Hz, 1H, H-3'), 7.46 (dd, $J = 8.6, 2.3$ Hz, 1H, H-5'), 6.90 (d, $J = 8.6$ Hz, 1H, H-6'). ^{13}C NMR (125 MHz, CDCl_3) δ : 143.6 (C-1), 144.4 (C-2), 111.9 (C-3), 119.7 (C-4), 110.8 (C-5), 130.1 (C-6), 151.4 (C-1'), 115.6 (C-2'), 136.4 (C-3'), 118.2 (C-4'), 132.1 (C-5'), 121.6 (C-6').

2-(2',4'-Dibromophenoxy)-3,5,6-tribromophenol (19): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 7.55 (s, 1H, H-4), 6.12 (br s, 1-H, 1-OH), 7.78 (d, $J = 2.3$ Hz, 1H, H-3'), 7.27 (dd, $J = 9.2, 2.3$ Hz, 1H, H-5'), 6.41 (d, $J = 9.2$ Hz, 1H, H-6'). ^{13}C NMR (125 MHz, CDCl_3) δ : 148.2 (C-1), 138.7 (C-2), 116.8 (C-3), 128.0 (C-4), 122.2 (C-5), 113.4 (C-6), 152.2 (C-1'), 112.7 (C-2'), 136.1 (C-3'), 115.8 (C-4'), 131.3 (C-5'), 115.9 (C-6').

3,6-Dibromo-2-(2',4'-dibromophenoxy) phenol (20): white powder. ^1H NMR (500 MHz, CDCl_3) and ^{13}C NMR (125 MHz, CDCl_3): see Table 3.1.

2-(2',4'-Dibromophenoxy)-4,6-dibromoanisole (21): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 6.92 (d, $J = 2.3$ Hz, 1H, H-4), 7.51 (d, $J = 2.3$ Hz, 1H, H-6), 3.91 (s, 3H, O-Me), 7.80 (d, $J = 2.3$ Hz, 1H, H-3'), 7.40 (dd, $J = 8.6, 2.3$ Hz, 1H, H-5'), 6.76 (d, $J = 8.6$ Hz, 1H, H-6'). ^{13}C NMR (125 MHz, CDCl_3) δ : 147.6 (C-1), 150.0 (C-2), 121.9 (C-3), 116.6 (C-4), 131.0 (C-5), 119.3 (C-6), 61.3 (O-Me), 152.1 (C-1'), 114.7 (C-2'), 136.2 (C-3'), 117.2 (C-4'), 131.9 (C-5'), 120.4 (C-6').

Cholesterol (22): white powder.

5 α , 8 α -Epidioxycholest-6-en-3 β -ol (23): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 1.93 (m, 2H, H-1), 1.82 (m, 2H, H-2), 3.95 (tt, $J = 11.5, 5.1$ Hz, 1H, H-3), 2.09 (m, 2H, H-4), 6.22 (d, $J = 8.5$ Hz, 1H, H-6), 6.49 (d, $J = 8.5$ Hz, 1H, H-7), 1.47 (m, 2H, H-9), 1.96 (m, 2H, H-12), 1.53 (m, 1H, H-14), 1.61 (m, 2H, H-15), 1.89 (m, 2H, H-16), 1.15 (m, 1H, H-17), 0.78 (s, 3H, H-18), 0.86 (s, 3H, H-19), 1.36 (m, 1H, H-20), 0.88 (d, $J = 6.5$ Hz, 3H, H-21), 1.32, 0.98 (m, 2H, H-22), 1.33, 1.13 (m, 2H, H-23), 1.09 (m, 2H, H-24), 1.49 (m, 1H, H-25), 0.84 (d, $J = 6.6$ Hz, 3H, H-26), 0.85 (d, $J = 6.6$ Hz, 3H, H-27).

Inseparable epidioxysterol (24a/b): white powder; ^1H NMR (500 MHz, CDCl_3) δ : 3.93 (m, 1H, H-3), 6.24 (d, $J = 8.6$ Hz, 1H, H-6), 6.50 (d, $J = 8.6$ Hz, 1H, H-7), 0.81 (s, 3H, H-18), 0.88 (s, 3H, H-19), 0.95 (d, $J = 7.5$ Hz, 3H, H-21), 0.86 and 0.87 (d, $J = 7.5$ Hz, 1.5H, H-26), 0.88 and 0.88 (d, $J = 7.5$ Hz, 1.5H, H-27), 5.73 and 5.73 (dd, $J = 17.8, 11.5$ Hz, 0.5H, H-28), 5.27 and 5.28 (dd, $J = 11.5, 1.1$ Hz, 0.5H, H-29) 5.15 (d, $J = 17.8, 1.1$ Hz, 1H, H-29).

4 Experimental detail of chapter 4

Marine sponge material

The *Gelliodes* sp. marine sponges were collected from Hon Thom, Phu Quoc, Vietnam, in April 2018, and positively identified by Quang Minh Thai at the Vietnam Academy of Science and Technology, Institute of Oceanography, Nha Trang (Vietnam). A voucher specimen (E:55739) was deposited with the Ocean-ography Museum, Institute of Oceanography in NhaTrang, Vietnam.

Extraction and isolation

Dried *Gelliodes* sp. (611 g) was macerated with MeOH (2 L $\times$ 3) for 2 h with sonication to afford the crude extract, obtained after filtration and removal of the solvent using a rotary evaporator. This crude extract (62.3 g) was triturated in water (50 mL) and successively partitioned with *n*-hexane to obtain the *n*-hexane (8.9 g) extract, after removal of the solvents using a rotatory evaporator. The crude extract subjected to HP-20 column chromatography, using MeOH and H₂O, to give the H₂O (containing mainly salts, not evaporated), 50% MeOH (1.1 g), and MeOH (2.5 g) fractions, respectively. The MeOH fraction was subjected to normal phase silica gel open column chromatography using solvent systems composed of CHCl₃/MeOH (9.9:0.1, 4.9:0.1, 9.5:0.5, 9:1, 8:2, 7:3, 6:4, 5:5, 4:6, 0:10) with increasing polarities. In total, 40 fractions of 300 mL each were collected and pooled into 10 combined fractions after TLC profiling.

The combined fraction 2 (1 g), obtained from CHCl₃/MeOH (4.9:0.1), was subjected to reverse phase column chromatography using solvent systems composed of CHCl₃/MeOH with increasing polarities to yield the *F*_{1–11} sub fractions. Compound **27**

(6.0 mg) was purified from sub fraction F_4 (41.4 mg) using preparative reverse phase TLC eluted with $H_2O/MeOH$ (4:1). Compound **29** (2.3 mg, t_R 22 min) was isolated from sub fraction F_6 (206 mg) using preparative reverse phase HPLC equipped 5C₁₈-AR- II (10 mm I.D $\times$ 250 mm) column with gradient elution by $H_2O/MeOH$ (1:4) to only MeOH (flow rate 2 mL/min, monitored at 254 nm, and running time 60 min). Compound **28** (2.4 mg) was obtained from sub fraction F_7 (20.2 mg) using preparative reverse phase TLC with $H_2O/MeOH$ (1:4) solvent system after Sephadex LH-20 column chromatography eluted with only MeOH. The combined fraction 3 (259.8 mg), obtained from $CHCl_3/MeOH$ (9.5:0.5) subjected reverse phase column chromatography with gradient elution by $H_2O/MeOH$ (1:1) to only MeOH and preparative normal phase TLC eluted with $CHCl_3/MeOH$ (20:1) to yield **25** (2.1 mg). The 50% MeOH fraction was subjected to normal phase silica gel open column chromatography using solvent systems composed of $CHCl_3/MeOH$ (9.9:0.1, 4.9:0.1, 9.5:0.5, 9:1, 8:2, 7:3, 6:4, 5:5, 4:6, 0:10) with increasing polarities. Compounds **26** (5.4 mg), **30** (2.2 mg), and **31** (1.0 mg) were purified from each fractions 2 (131.9 mg), 3–4 (33.2, 37.6 mg), and 6 (62.8 mg) using preparative reverse phase HPLC equipped 5C₁₈-AR- II (10 mm I.D $\times$ 250 mm) column with gradient elution by $H_2O/MeOH$ (2:3) to only MeOH.

Gelliodesinic acid (25): colorless oil. IR (KBr) ν_{max} : 3419, 2925, 2854, 1756, 1712, 1654, 1454, 1383, 1256, 1100 cm^{-1} . UV (MeOH) λ_{max} (log ϵ): 206 (4.1), 212 (4.3), 218 (4.4), 228 (4.3), 294 (3.3). 1H NMR (500 MHz, $CDCl_3$) and ^{13}C NMR (125 MHz, $CDCl_3$): see Table 4.2. HR-ESI-MS (+): m/z 221.1151 $[M+Na]^+$ (calcd. for C₁₁H₁₉O₃Na, 221.1154).

3-(Propan-2-ylidene)indolin-2-one (26): yellowish powder; 1H NMR (400

MHz, CDCl₃) δ : 7.52 (d, J = 7.7 Hz, 2H, H-4), 7.19 (td, J = 7.7, 1.0 Hz, 2H, H-5), 7.02 (td, J = 7.7, 1.0 Hz, 2H, H-6), 6.84 (d, J = 7.7 Hz, 2H, H-7), 2.61 (s, 3H, H-10), 2.38 (s, 3H, H-9). ¹³C NMR (100 MHz, CDCl₃) δ : 169.0 (C-2), 155.4 (C-8), 138.8 (C-3a), 127.2 (C-5), 124.0 (C-7a), 123.4 (C-7), 122.5 (C-3), 121.3 (C-6), 108.8 (C-7), 25.0 (C-9), 22.8 (C-10). ESI-MS (+): m/z 174.1 ([M+H]⁺, 100), 196.1 ([M+Na]⁺, 10).

Sarcotin H (27): yellowish oil; ¹H NMR (500 MHz, CD₃OD) δ : 7.36 (br s, 1H, H-1), 6.27 (br s, 1H, H-2), 7.30 (br s, 1H, H-3), 3.04 (s, 2H, H-5), 6.85 (s, 1H, H-7), 2.15 (m, 2H, H-10), 1.49 (m, 2H, H-11), 1.95 (m, 2H, H-12), 1.64 (s, 3H, H-14), 5.15 (t, J = 6.8 Hz, 1H, H-15), 1.97 (m, 2H, H-16), 1.40, 1.47 (m, 2H, H-17), 2.73 (m, 1H, H-18), 1.05 (d, J = 6.8 Hz, 3H, H-19), 5.27 (d, J = 10.0 Hz, 1H, H-20), 1.72 (s, 3H, H-25) 3.21 (s, 3H, OCH₃).

(7*E*, 12*E*, 20*Z*, 18*S*)-Variabilin (28): yellowish oil; ¹H NMR (500 MHz, CDCl₃) δ : 7.34 (br s, 1H, H-1), 6.28 (br s, 1H, H-2), 7.21 (br s, 1H, H-4), 2.42 (t, J = 7.6 Hz, 2H, H-5), 2.23 (q, J = 7.2 Hz, 2H, H-6), 5.17 (t, J = 7.0 Hz, 1H, H-7), 1.59 (s, 3H, H-9), 1.94 (m, 4H, H-10 and H-15), 2.06 (m, 2H, H-11), 5.08 (t, J = 6.9 Hz, 1H, H-12), 1.56 (s, 3H, H-14), 1.26 (m, 4H, H-16 and H-17), 2.79 (br s, 1H, H-18), 1.05 (d, J = 6.8 Hz, 3H, H-19), 5.26 (d, J = 10.0 Hz, 1H, H-20), 1.81 (s, 3H, H-25).

Sulawesin C (29): yellowish oil; ¹H NMR (400 MHz, CD₃OD) δ : 7.38 (br s, 1H, H-1), 6.29 (br s, 1H, H-2), 7.28 (br s, 1H, H-4), 3.70 (br s, 2H, H-5), 5.88 (br s, 1H, H-7), 2.30 (t, J = 7.4 Hz, 2H, H-10), 2.11 (d, J = 7.0 Hz, 2H, H-11), 5.05 (t, J = 7.3 Hz, 1H, H-12), 1.45 (s, 3H, H-14), 1.92 (m, 2H, H-15), 1.38 (m, 2H, H-16), 1.28, 1.33 (m, 2H, H-

17), 2.70 (m, 1H, H-18), 1.02 (d, 6.8 Hz, 3H, H-19), 5.23 (d, $J = 10$ Hz, 1H, H-20), 1.65 (s, 3H, H-25), 7.40 (br s, 1H, H-1'), 6.34 (br s, 1H, H-2'), 7.23 (br s, 1H, H-4'), 5.23 (s, 1H, H-5'), 5.86 (br s, 1H, H-7'), 7.11 (br s, H-9'), 2.35 (t, $J = 7.4$ Hz, 2H, H-10'), 2.18 (q, $J = 7.4$ Hz, 2H, H-11'), 5.11 (t, $J = 7.3$ Hz, 1H, H-12'), 1.50 (s, 3H, H-14'), 1.92 (m, 2H, H-15'), 1.38 (m, 2H, H-16'), 1.28, 1.33 (m, 2H, H-17'), 2.70 (m, 1H, H-18'), 1.02 (d, $J = 6.8$ Hz, 3H, H-19'), 5.23 (d, $J = 10$ Hz, 1H, H-20'), 1.65 (s, 3H, H-25').

1H-Indole-3-carboxylic acid (30): yellowish powder; ^1H NMR (500 MHz, DMSO- d_6) δ : 7.96 (s, 1H, H-2), 7.45 (d, $J = 7.1$ Hz, 1H, H-4), 7.15 (m, 2H, H-5 and H-6), 8.02 (d, $J = 7.7$ Hz, 1H, H-7).

3-(Hydroxyacetyl)indole (31): yellowish powder; ^1H NMR (500 MHz, CDCl_3) δ : 7.91 (d, $J = 3.1$ Hz, 1H, H-2), 8.26 (m, 1H, H-4), 7.32 (m, 2H, H-5 and H-6), 7.44 (m, 1H, H-7), 4.71 (s, 2H, H-9).

References

1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A (2018) Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68:394–424.
2. Graidist P, Martla M, Sukpondma Y (2015) Cytotoxic activity of *Piper cubeba* extract in breast cancer cell lines. *Nutrients* 7:2707–2718.
3. Limbo CA, Jacinto SD (2019) Cytotoxic potential and phytochemical profile of extracts from *Garcinia rubra* Merr. leaves. *Cancer Research* 15:38–46.
4. Koehn FE, Carter GT (2005) The evolving role of natural products in drug discovery. *Nat Rev Drug Discov* 4:206–620.
5. Walsh V, Goodman J (1999) Cancer chemotherapy, biodiversity, public and private property: the case of the anti-cancer drug taxol. *Soc Sci Med* 49:1215–1225.
6. Mugera GM (1977) Useful drugs and cancer causing chemicals in Kenya medical and toxic plants; department of veterinary pathology and microbiology: Nairobi, Kenya.
7. Bourhill T, Narendran A, Johnston R.N (2017) Enzastaurin A: lesson in drug development. *Crit Rev Onco Hematol* 112:72–79.
8. Wang YQ, Miao ZH (2013) Marine-derived angiogenesis inhibitors for cancer therapy. *Mar Drugs* 11:903–933.
9. Ritchie JWA; Williams RJ (2015) Cancer research UK centre for drug development: translating 21st-century science into the cancer medicines of tomorrow. *Drug Discov Today* 20:995–1003.

10. Khalifa SAM, Elias N, Farag MA, Chen L, Saeed A, Hegazy MF, Moustafa MS, Abd El-Wahed A, Al-Mousawi SM, Musharraf SG, Chang FR, Iwasaki A, Suenaga K, Alajlani M, Göransson U, El-Seedi HR. (2019) Marine natural products: a source of novel anticancer drugs. *Mar Drugs* 17: pii: E491.
11. Coseri S (2009) Natural products and their analogues as efficient anticancer drugs. *Mini Rev Med Chem* 9:560–571.
12. Newman DJ (2008) Natural products as leads to potential drugs: an old process or the new hope for drug discovery? *J Med Chem* 51:2589–2599.
13. Newman DJ, Gragg GM, Snader KM (2003) Natural products as sources of new drugs over the period 1981–2002. *J Nat Prod* 66:1022–1037.
14. Hentschel U, Schmid M, Wagner M, Fieseler L, Gernert C, Hacker J (2001) Isolation and phylogenetic analysis of bacteria with antimicrobial activities from the Mediterranean sponges *Aplysina aerophoba* and *Aplysina cavernicola*. *FEMS Microbiol Ecol* 35:305–312.
15. Thomas TRA, Kavlekar DP, LokaBharathi PA (2010) Marine drugs from sponge-microbe association—a review. *Mar Drugs* 8:1417–1468.
16. Radjasa OK, Sabdono A, Junaidi J, Zocchi E (2007) Richness of secondary metabolite-producing marine bacteria associated with sponge *Haliclona* sp.. *Int J Pharmacol* 3:275–279.
17. Mehbub MF, Lei J, Franco C, Zhang W (2014) Marine sponge derived natural products between 2001 and 2010: trends and opportunities for discovery of bioactives. *Mar Drugs* 12:4539–4577.
18. Bergmann W, Feeney R.J (1951) Contributions to the study of marine products. XXXII. The nucleosides of sponges. I. *J Org Chem* 16:981–987.

19. Bergmann W, Feeney RJ (1950) The isolation of a new thymine pentoside from sponges 1. *J Am Chem Soc* 72:2809–2810.
20. Faulkner DJ (2000) Marine natural products. *Nat Prod Rep* 17:7–55.
21. Faulkner DJ (2002) Marine natural products. *Nat Prod Rep* 19:1–48.
22. Rajendran S. (2019) Marine sponges: repositories of bioactive compounds with medicinal applications. *Int J Chemtech Res* 12:26–48.
23. Perdicaris S, Vlachogianni T, Valavanidis A. (2013) Bioactive natural substances from marine sponges: new developments and prospects for future pharmaceuticals. *Nat Prod Chem Res* 1:2329–6836.
24. Anjum K, Abbas SQ, Shah SA, Akhter N, Batool S, Hassan SS (2016) Marine sponges as a drug treasure. *Biomol Ther* 24:347–362.
25. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2010) Marine natural products. *Nat Prod Rep* 27:165–237.
26. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2011) Marine natural products. *Nat Prod Rep* 28:196–268.
27. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2012) Marine natural products. *Nat Prod Rep* 29:144–222.
28. Blunt JW, Copp BR, Keyzers RA, Munro MHG, Prinsep MR (2013) Marine natural products. *Nat Prod Rep* 30:237–323.
29. Blunt JW, Copp BR, Keyzers RA, Munro MHG, Prinsep MR (2014) Marine natural products. *Nat Prod Rep* 31:160–258.
30. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2003) Marine natural products. *Nat Prod Rep* 20:1–48.
31. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2004) Marine

- natural products. *Nat Prod Rep* 21:1–49.
32. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2005) Marine natural products. *Nat Prod Rep* 22:15–61.
 33. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2006) Marine natural products. *Nat Prod Rep* 23:26–78.
 34. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2007) Marine natural products. *Nat Prod Rep* 24:31–86.
 35. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2008) Marine natural products. *Nat Prod Rep* 25:35–94.
 36. Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR (2009) Marine natural products. *Nat Prod Rep* 26:170–244.
 37. Shin J, Rho JR, Seo Y, Lee HS, Cho KW, Sim CJ (2001) Sarcotragins A and B, new sesterterpenoid alkaloids from the sponge *Sarcotragus* sp.. *Tetrahedron Lett* 42:3005–3007.
 38. Cichewicz RH, Valeriote FA, Crews P (2004) Psymberin, a potent sponge-derived cytotoxin from *Psammocinia* distantly related to the pederin family. *Org Lett* 6:1951–1954.
 39. Johnson TA, Amagata T, Oliver AG, Tenney K, Valeriote FA, Crews P (2008) The unexpected isolation of CTP-431, a novel thiopyrone from the sponge *Cacospongia mycofijiensis*. *J Org Chem* 73:7255–7259.
 40. Bae JM, Jeon JE, Lee YJ, Lee HS, Sim CJ, Oh KB, Shin JH (2011) Sesterterpenes from the tropical sponge *Coscinoderma* sp.. *J Nat Prod* 74:1805–1811.
 41. Ovenden SPB, Nielson JL, Liptrot CH, Willis RH, Tapiolas DM, Wright AD, Motti CA (2011) Sesquiterpene benzoxazoles and sesquiterpene quinones from

- the marine sponge *Dactylospongia elegans*. *J Nat Prod* 74:65–68.
42. Prawat H, Mahidol C, Kaweetripob W, Wittayalai S, Ruchirawat S (2012) Iodosesquiterpene hydroquinone and brominated indole alkaloids from the Thai sponge *Smenospongia* sp.. *Tetrahedron* 68:6881–6886.
 43. Davis RA, Buchanan MS, Duffy S, Avery VM, Charman SA, Charman WN, White KL, Shackleford DM, Edstein MD, Andrews KT (2012) Antimalarial activity of pyrroloiminoquinones from the Australian marine sponge *Zyzzya* sp.. *J Med Chem* 55:5851–5858.
 44. Jeon J, Na Z, Jung M, Lee, HS, Sim CJ, Nahm K, Oh KB, Shin J (2010) Discorhabdins from the Korean marine sponge *Sceptrella* sp.. *J Nat Prod* 73:258–262.
 45. Wang W, Lee Y, Lee TG, Mun B, Giri AG, Lee J, Kim H, Hahn D, Yang I. Chin J (2012) Phorone A and isophorbasone A, sesterterpenoids isolated from the marine sponge *Phorbas* sp.. *Org Lett* 14:4486–4489.
 46. Chao CH, Chou KJ, Wang GH, Wu YC, Wang LH, Chen JP, Sheu JH, Sung PJ (2010) Nortriterpenoids and related peroxides from the Formosan marine sponge *Negombata corticata*. *J Nat Prod* 73:1538–1543.
 47. Rogers EW, Molinski TF (2005) A cytotoxic carotenoid from the marine sponge *Prianos osiros*. *J Nat Prod* 68:450–452.
 48. Kura K, Kubota T, Fromont J, Kobayashi J (2011) Pyrinodemins E and F, new 3-alkylpyridine alkaloids from sponge *Amphimedon* sp.. *Bioorg Med Chem Lett* 21:267–270.
 49. Hitora Y, Takada K, Okada S, Ise Y, Matsunaga S (2011) (–)-Duryne and its homologues, cytotoxic acetylenes from a marine sponge *Petrosia* sp.. *J Nat Prod*

74:1262–1267.

50. Umeyama A, Matsuoka N, Mine R, Nakata A, Arimoto E, Matsui M, Shoji N, Arihara S, Takei M, Hashimoto T. (2010) Polyacetylene diols with antiproliferative and driving Th1 polarization effects from the marine sponge *Callyspongia* sp.. *J Nat Med* 64:93–97.
51. Rotem M, Kashman Y (1979) New polyacetylenes from the sponge *Siphonochalina* sp.. *Tetrahedron Lett* 34:3193–3196.
52. Hooper JNA, van Soest RW (2002) 'Systema Porifera: a guide to the classification of sponges', Springer US, New York.
53. Carmely S, Kashman Y (1986) The study of sipholanes by two-dimensional NMR spectroscopy. *Magn Reson Chem* 24:332–336.
54. Jain S, Laphookhieo S, Shi Z, Fu LW, Akiyama S, Chen ZS, Youssef DT, van Soest RW, El Sayed KA (2007) Reversal of p-glycoprotein-mediated multidrug resistance by sipholane triterpenoids. *J Nat Prod* 70:928–931.
55. Kashman Y, Yosief T, Carmeli S (2001) New triterpenoids from the Red Sea sponge *Siphonochalina siphonella*. *J Nat Prod* 64:175–180.
56. El-Beih AA, El-Desoky AH, Al-Hammady MA, Elshamy AI, Hegazy MF, Kato H, Tsukamoto S (2018) New inhibitors of RANKL-induced osteoclastogenesis from the marine sponge *Siphonochalina siphonella*. *Fitoterapia* 128:43–49.
57. Ali MS, Amina M, Al-Lohedan HA, Al Musayeib NM (2017) Elucidation of the interaction of human serum albumin with anti-cancer sipholane triterpenoid from the Red Sea sponge. *Luminescence* 32: 223–230.
58. Amina M, Ali MS, Al-Musayeib NM, Al-Lohedan HA (2016) Biophysical

- characterization of the interaction of bovine serum albumin with anticancer sipholane triterpenoid from the Red Sea sponge. *J Mol Liq* 220:931–938.
59. Al-Massarani SM, El-Gamal AA, Al-Said MS, Al-Lihaibi SS, Basoudan OA (2015) In vitro cytotoxic, antibacterial and antiviral activities of triterpenes from the Red Sea sponge, *Siphonochalina siphonella*. *Trop J Pharm Res* 14:33–40.
 60. Angawi RF, Saqer E, Abdel-Lateff A, Badria FA, Ayyad SE (2014) Cytotoxic neviotane triterpene-type from the Red Sea sponge *Siphonochalina siphonella*. *Pharmacogn Mag* 10:s334–s341.
 61. Ayyad SE, Angawy R, Alarif WM, Saqer EA, Badria FA (2014) Cytotoxic polyacetylenes from the Red Sea sponge *Siphonochalina siphonella*. *Z Naturforsch* 69c:117–123.
 62. Sobahi TRA, Ayyad SN, Abdel-Lateff A, Algandaby MM, Alorfi HS, Abdel-Naim AB. (2017) Cytotoxic metabolites from *Callyspongia siphonella* display antiproliferative activity by inducing apoptosis in HCT-116 cells. *Pharmacogn Mag* 13:S37–S40.
 63. Youssef DT, van Soest RW, Fusetani N (2003) Callyspongamide A, a new cytotoxic polyacetylenic amide from the Red Sea sponge *Callyspongia fistularis*. *J Nat Prod* 66:861–862.
 64. Youssef DT, van Soest RW, Fusetani N (2003) Callyspongenols A–C, new cytotoxic C22-polyacetylenic alcohols from a Red Sea sponge, *Callyspongia* species. *J Nat Prod* 66:679–681.
 65. Mejia EJ, Magranet LB, De Voogd NJ, Tendyke K, Qiu D, Shen YY, Zhou Z, Crews P (2013) Structures and cytotoxic evaluation of new and known acyclic ene-yne from an American Samoa *Petrosia* sp. sponge. *J Nat Prod* 76:425–432.

66. Grundmann F, Dill V, Dowling A, Thanwisai A, Bode E, Chantratita N, Ffrench-Constant R, Bode HB (2012) Identification and isolation of insecticidal oxazoles from *Pseudomonas* spp.. *Beilstein J Org Chem* 8:749–752.
67. Breitmaier E, Voelter W (1987) In carbon-13 NMR spectroscopy, VCH Verlagsgesellschaft: Weinheim, pp.192–194.
68. Trindade-Silva AE, Rua C, Silva GG, Dutilh BE, Moreira AP, Edwards RA, Hajdu E, Lobo-Hajdu G, Vasconcelos AT, Berlinck RG, Thompson FL (2012) Taxonomic and functional microbial signatures of the endemic marine sponge *Arenosclera brasiliensis*. *PLoS One* 7:e39905.
69. Zhang D, Sun W, Feng G, Zhang F, Anbuezhian R, Li Z, Jiang Q (2015) Phylogenetic diversity of sulphate-reducing desulfovibrio associated with three South China Sea sponges. *Lett Appl Microbiol* 60:504–512.
70. Torres YR, Berlinck RG, Magalhães A, Schefer AB, Ferreira AG, Hajdu E, Muricy G (2000) Arenosclerins A–C and haliclonacyclamine E, new tetracyclic alkaloids from a Brazilian endemic Haplosclerid sponge *Arenosclera brasiliensis*. *J Nat Prod* 63:1098–1105.
71. Torres YR, Berlinck RG, Nascimento GG, Fortier SC, Pessoa C, de Moraes MO (2002) Antibacterial activity against resistant bacteria and cytotoxicity of four alkaloid toxins isolated from the marine sponge *Arenosclera brasiliensis*. *Toxicon* 40:885–891.
72. Shridhar DM, Mahajan GB, Kamat VP, Naik CG, Parab RR, Thakur NR, Mishra PD (2009) Antibacterial activity of 2-(2',4'-dibromophenoxy)-4,6-dibromophenol from *Dysidea granulosa*. *Mar Drugs* 7:464–471.
73. Yamaoka Y, Ohta S, Carmona ML, Oclarit JM (2006) Content and composition

- of brominated compounds in marine sponges. *Bull. Soc. Sea Water Sci* 60:195–198.
74. Marsh G, Stenutz R, Bergman Å (2003) Synthesis of hydroxylated and methoxylated polybrominated diphenyl ethers—natural products and potential polybrominated diphenyl ether metabolites. *Eur J Org Chem* 14:2566–2576.
 75. Calcul L, Chow R, Oliver AG, Tenney K, White KN, Wood AW, Fiorilla C, Crews P (2009) NMR strategy for unraveling structures of bioactive sponge-derived oxy-polyhalogenated diphenyl ethers. *J Nat Prod* 72:443–449.
 76. Zhang H, Skildum A, Stromquist E, Rose-Hellekant T, Chang LC (2008) Bioactive polybrominated diphenyl ethers from the marine sponge *Dysidea* sp.. *J Nat Prod* 71:262–264.
 77. Cameron GM, Stapleton BL, Simonsen SM, Brecknell DJ, Garson MJ (2000) New sesquiterpene and brominated metabolites from the tropical marine sponge *Dysidea* sp.. *Tetrahedron* 56:5247–5252.
 78. Kovganko NV, Kashkan ZN, Borisov EV (2000) ^{13}C NMR spectra of functionally substituted 3 β -chloroderivatives of cholesterol and β -sitosterol. *Chem Nat Compd* 36:595–598.
 79. Liu Y, Yan H, Wen K, Zhang J, Xu T, Wang L, ZHOU X, Yang X (2011) Identification of epidioxysterol from South China Sea urchin *Tripneustes gratilla* Linnaeus and its cytotoxic activity. *J Food Biochem* 35:932–938.
 80. Sera Y, Adachi K, Shizuri Y (1999) A new epidioxy sterol as an antifouling substance from a Palauan marine sponge, *lendenfeldia chondrodes*. *J Nat Prod* 62:152–154.
 81. Núñez Pons L, Calcinai B, Gates RD. (2017) Who's there?- First morphological

- and DNA barcoding catalogue of the shallow Hawai'ian sponge fauna. *PLoS One* 12:e0189357.
82. Tanaka Y and Inoue T, (1986) A new aldehydic carotenoid gelliodesxanthin from sea sponge *Gelliodes callista*. *Nippon Suisan Gakk* 53:1271–1273.
 83. Zhen X, Gong T, Liu F, Zhang PC, Zhou WQ, Li Y, Zhu P. (2015) A new analogue of echinomycin and a new cyclic dipeptide from a marine-derived *Streptomyces* sp. LS298. *Mar Drugs* 13:6947–6961.
 84. Gong T, Zhen X, Li BJ, Yang JL, Zhu P. (2015) Two new monoterpenoid α -pyrones from a fungus *Nectria* sp. HLS206 associated with the marine sponge *Gelliodes carnosa*. *J Asian Nat Prod Res* 17:633–637.
 85. Zhu H, Hua XX, Gong T, Pang J, Hou Q, Zhu P. (2013) Hypocreaterpenes A and B, cadinane-type sesquiterpenes from a marine-derived fungus, *Hypocreales* sp.. *Phytochem Lett* 6:392–396.
 86. Harrowven DC, Stenning KJ, Whiting S, Thompson T, Walton R. (2011) CH activation and CH₂ double activation of indolines by radical translocation: understanding the chemistry of the indoliny radical. *Org Biomol Chem*. 9:4882–4885.
 87. Liu Y, Hong J, Lee CO, Im KS, Kim ND, Choi JS, Jung JH (2002) Cytotoxic pyrrolo- and furanoterpenoids from the sponge *Sarcotragus* species. *J Nat Prod* 65:1307–1314.
 88. Abdjul DB, Yamazaki H, Kanno SI, Wewengkang DS, Rotinsulu H, Sumilat DA, Ukai K, Kapojos MM, Namikoshi M (2017) Furanoterpenes, new types of protein tyrosine phosphatase 1B inhibitors, from two Indonesian marine sponges, *Ircinia* and *Spongia* spp.. *Bioorg Med Chem Lett* 27:1159–1161.

89. Afifi AH, Kagiya I, El-Desoky AH, Kato H, Mangindaan REP, de Voogd NJ, Ammar NM, Hifnawy MS, Tsukamoto S (2017) Sulawesins A–C, furanosesterterpene tetronic acids that inhibit USP7, from a *Psammocinia* sp. marine sponge. *J Nat Prod* 80:2045–2050.
90. Yan X, Xue-ping S, Xiu-guo Z, Wan-ping T (2015) Insecticidal activity of compounds from the South China Sea invertebrate *Scapharca subcrenata*. *Chem Nat Compd* 51:800–802.
91. Yang SW, Cordell GA. (1997) Metabolism studies of indole derivatives using a staurosporine producer, *Streptomyces staurosporeus*. *J Nat Prod* 60:44–48.
92. Emura C, Higuchi R, Miyamoto T (2010) Synthetic studies on the natural multidrug resistance modulator, irciniasulfonic acid B. *Chem Lett* 39:1002–1003.
93. Phan VK, Duong DT, Do TT, Tran QH, Nguyen NT, Tran HH, Hoang Le AT, Pham YH, DO TT, Nguyen NX, Bui TH, Oh H, Kim YC, Chau MV (2017) Constituents from *Ircinia echinata* and their antiproliferative effect on six human cancer cell strains. *Lett Org Chem* 14:248–253.
94. Tan LT, Okino T, Gerwick WH (2000) Hermitamides A and B, toxic malyngamide-type natural products from the marine cyanobacterium *Lyngbya majuscula*. *J Nat Prod* 63: 952–955.
95. Agarwal V, Blanton JM, Podell S, Taton A, Schorn MA, Busch J, Lin Z, Schmidt EW, Jensen PR, Paul VJ, Biggs JS, Golden JW, Allen EE, Moore BS (2017) Metagenomic discovery of polybrominated diphenyl ether biosynthesis by marine sponges. *Nat Chem Biol* 13: 537–543.
96. Matsunaga H, Katano M, Yamamoto H, Fujito H, Mori M, Takata K (1990) Cytotoxic activity of polyacetylene compounds in *Panax ginseng* CA Meyer.

Chem Pharm Bull 38: 3480–3482.

97. Zhong YF, Wang LL, Yin LL, An J, Hou ML, Zheng KW, Zhang XY, Wu MH, Yu ZQ, Sheng GY, Fu JM (2011) Cytotoxic effects and oxidative stress response of six PBDE metabolites on human L02 cells. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 46: 1320–1327.

List of publications

This doctoral thesis summarizes the full contents of the following publications.

1. Ki DW, Awouafack MD, Wong CP, Nguyen HM, Thai QM, Ton Nu LH, Morita H (2019) Brominated diphenyl ethers including a new tribromiododiphenyl ether from the Vietnamese marine sponge *Arenosclera* sp. and their antibacterial activities. *Chem Biodivers* 16:e1800593.
2. Ki DW, El-Desoky AH, Wong CP, Ghani MA, El-Beih AA, Mineyuki M, Morita H (2020) New cytotoxic polyacetylene alcohols from the Egyptian marine sponge *Siphonochalina siphonella*. *J Nat Med* 74:409–414.
3. Ki DW, El-Desoky AH, Kodama T, Wong CP, Ghani MA, El-Beih AA, Mineyuki M, Morita H (2020) New cytotoxic polyacetylene amides from the Egyptian marine sponge *Siphonochalina siphonella*. *Fitoterapia* 142:104511.
4. Ki DW, Kodama T, El-Desoky AH, Wong CP, Nguyen HM, Thai QM, Ton Nu LH, Morita H (2020) Chemical constituents of the Vietnamese marine sponge *Gelliodes* sp. and their cytotoxic activities. *Chem Biodivers* accepted <https://doi.org/10.1002/cbdv.202000303>.

Acknowledgements

I would like to express my deepest and sincere gratitude to my supervisor Prof. Hiroyuki Morita, Section of Natural Products Chemistry & Drug Discovery, Institute of Natural Medicine, University of Toyama, Japan, for his valuable instruction, guidance, suggestion and encouragements on my research to achieve my PhD degree.

I am also greatly indebted to Assistant Prof. Takeshi Kodama, Section of Natural Products Chemistry & Drug Discovery, Institute of Natural Medicine, University of Toyama, Japan, for his valuable suggestions, support, and comments throughout this study.

I am highly grateful to former Assistant Prof. Chin Piow Wong, for his guidance and kindness to my study.

I would like to express of my sincere gratitude to Prof. Kastuko Komatsu, Section of Pharmacognosy, Institute of Natural Medicine, University of Toyama, Japan, and Prof. Yuji Matsuya, Laboratory of Synthetic and Medicinal Chemistry, Department of Pharmaceutical Sciences, University of Toyama, Japan, for their comments, suggestions and evaluating my thesis.

I am also greatly indebted to Prof. Ahmed A. El-Beih and Dr. Ahmed H. El-Desoky, Department of Pharmacognosy, National Research Center, Egypt, and Associate Prof. Maurice Ducret Awouafack, Natural Products Chemistry Research Unit, Dschang University, Cameroon, for their valuable suggestions, support, and comments throughout this study.

I would like to thanks to express my thanks to Prof. Mineyuki Mizuguchi, Laboratory of Structural Biology, Department of Pharmaceutical Sciences, University of Toyama, Japan, for the help to measure NMR.

I extend my heartfelt thanks to all members in Section of Natural Products Chemistry & Drug Discovery for the support and friendship during my study in the University of Toyama, Japan.

Finally, I am greatly indebted to my family for their love, understanding, support, and encouragement throughout the whole of my life.

Dae-Won Ki

Toyama, Japan

■ Supplementary data

List of contents

Figure S1. ^1H NMR spectrum (400 MHz) of siphonellamide A (**1**) in CDCl_3

Figure S2. ^{13}C NMR spectrum (100 MHz) of siphonellamide A (**1**) in CDCl_3

Figure S3. HMQC spectrum of siphonellamide A (**1**)

Figure S4. HMBC spectrum of siphonellamide A (**1**)

Figure S5. ^1H – ^1H COSY spectrum of siphonellamide A (**1**)

Figure S6. ^1H – ^1H TOCSY spectrum of siphonellamide A (**1**)

Figure S7. HR-ESI-MS spectrum of siphonellamide A (**1**)

Figure S8. ^1H NMR spectrum (400 MHz) of siphonellamide B (**2**) in CDCl_3

Figure S9. ^{13}C NMR spectrum (100 MHz) of siphonellamide B (**2**) in CDCl_3

Figure S10. HMQC spectrum of siphonellamide B (**2**)

Figure S11. HMBC spectrum of siphonellamide B (**2**)

Figure S12. ^1H – ^1H COSY spectrum of siphonellamide B (**2**)

Figure S13. ^1H – ^1H TOCSY spectrum of siphonellamide B (**2**)

Figure S14. HR-ESI-MS spectrum of siphonellamide B (**2**)

Figure S15. ^1H NMR spectrum (500 MHz) of siphonellamide C (**3**) in CDCl_3

Figure S16. HMQC spectrum of siphonellamide C (**3**)

Figure S17. HMBC spectrum of siphonellamide C (**3**)

Figure S18. ^1H – ^1H COSY spectrum of siphonellamide C (**3**)

Figure S19. ^1H – ^1H TOCSY spectrum of siphonellamide C (**3**)

Figure S20. HR-ESI-MS spectrum of Siphonellamide C (**3**)

Figure S21. ^1H NMR spectrum (500 MHz) of siphonellamide D (**4**) in CDCl_3

Figure S22. HMQC spectrum of siphonellamide D (**4**)

Figure S23. HMBC spectrum of siphonellamide D (**4**)

Figure S24. ^1H – ^1H COSY spectrum of siphonellamide D (**4**)

Figure S25. ^1H – ^1H TOCSY spectrum of siphonellamide D (**4**)

Figure S26. HR-ESI-MS spectrum of siphonellamide D (**4**)

Figure S27. ^1H NMR spectrum (400 MHz) of siphonellamide E (**5**) in CDCl_3

Figure S28. ^{13}C NMR spectrum (100 MHz) of siphonellamide E (**5**) in CDCl_3

Figure S29. HMQC spectrum of siphonellamide E (**5**)

Figure S30. HMBC spectrum of siphonellamide E (**5**)

Figure S31. ^1H – ^1H COSY spectrum of siphonellamide E (**5**)

Figure S32. ^1H – ^1H TOCSY spectrum of siphonellamide E (**5**)

Figure S33. HR-ESI-MS spectrum of siphonellamide E (**5**)

Figure S34. Ozonolysis of siphonellamide E (**5**) **A.** ESI-MS of ozonolysis products
B. ESI-MS/MS of m/z 301 $[\text{M}+\text{H}]^+$

Figure S35. ^1H NMR spectrum (500 MHz) of siphonellanol A (**6**) in CDCl_3

Figure S36. ^{13}C NMR spectrum (125 MHz) of siphonellanol A (**6**) in CDCl_3

Figure S37. HMQC spectrum of siphonellanol A (**6**)

Figure S38. HMBC spectrum of siphonellanol A (**6**)

Figure S39. ^1H – ^1H COSY spectrum of siphonellanol A (**6**)

Figure S40. ^1H – ^1H TOCSY spectrum of siphonellanol A (**6**)

Figure S41. HR-ESI-MS spectrum of siphonellanol A (**6**)

Figure S42. ^1H NMR spectrum (500 MHz) of siphonellanol B (**7**) in CDCl_3

Figure S43. ^{13}C NMR spectrum (100 MHz) of siphonellanol B (**7**) in CDCl_3

Figure S44. HMQC spectrum of siphonellanol B (**7**)

Figure S45. HMBC spectrum of siphonellanol B (**7**)

Figure S46. ^1H – ^1H COSY spectrum of siphonellanol B (**7**)

Figure S47. ^1H – ^1H TOCSY spectrum of siphonellanol B (**7**)

Figure S48. HR-ESI-MS spectrum of siphonellanol B (**7**)

Figure S49. ^1H NMR spectrum (400 MHz) of siphonellanol C (**8**) in CDCl_3

Figure S50. ^{13}C NMR spectrum (100 MHz) of Siphonellanol C (**8**) in CDCl_3

Figure S51. HMQC spectrum of siphonellanol C (**8**)

Figure S52. HMBC spectrum of siphonellanol C (**8**)

Figure S53. ^1H – ^1H COSY spectrum of siphonellanol C (**8**)

Figure S54. ^1H – ^1H TOCSY spectrum of siphonellanol C (**8**)

Figure S55. HR-ESI-MS spectrum of siphonellanol C (**8**)

Figure S56. ^1H NMR spectrum (500 MHz) of tribromiododiphenyl ether (**13**) in CDCl_3

Figure S57. ^{13}C NMR spectrum (125 MHz) of tribromiododiphenyl ether (**13**) in CDCl_3

Figure S58. HMQC spectrum of tribromiododiphenyl ether (**13**)

Figure S59. HMBC spectrum of tribromiododiphenyl ether (**13**)

Figure S60. ^1H – ^1H COSY spectrum of tribromiododiphenyl ether (**13**)

Figure S61. HR-ESI-MS spectrum of tribromiododiphenyl ether (**13**)

Figure S62. ^1H NMR spectrum (500 MHz) of gelliodesinic acid (**25**) in CDCl_3

Figure S63. ^{13}C NMR spectrum (125 MHz) of gelliodesinic acid (**25**) in CDCl_3

Figure S64. HMQC spectrum of gelliodesinic acid (**25**)

Figure S65. HMBC spectrum of gelliodesinic acid (**25**)

Figure S66. ^1H – ^1H COSY spectrum of gelliodesinic acid (**25**)

Figure S67. NOESY spectrum of gelliodesinic acid (**25**)

Figure S68. HR-ESI-MS spectrum of gelliodesinic acid (**25**)

Figure S69. ^1H NMR spectrum (400 MHz) of 3-(propan-2-ylidene)indolin-2-one (**26**) in CDCl_3

Figure S70. ^{13}C NMR spectrum (100 MHz) of 3-(propan-2-ylidene)indolin-2-one (**26**) in CDCl_3

Figure S71. HMQC spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)

Figure S72. HMBC spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)

Figure S73. ^1H – ^1H COSY spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)

Figure S74. ESI-MS spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)

Figure S1. ¹H NMR spectrum (400 MHz) of siphonellamide A (1) in CDCl₃

Figure S2. ¹³C NMR spectrum (100 MHz) of siphonellamide A (1) in CDCl₃

Figure S3. HMQC spectrum of siphonellamide A (**1**)

Figure S4. HMBC spectrum of siphonellamide A (**1**)

Figure S5. ^1H – ^1H COSY spectrum of siphonellamide A (**1**)

Figure S6. ^1H – ^1H TOCSY spectrum of siphonellamide A (**1**)

Figure S7. HR-ESI-MS spectrum of siphonellamide A (**1**)

Figure S8. ¹H NMR spectrum (400 MHz) of siphonellamide B (**2**) in CDCl₃

Figure S9. ^{13}C NMR spectrum (100 MHz) of siphonellamide B (**2**) in CDCl_3

Figure S10. HMQC spectrum of siphonellamide B (**2**)

Figure S11. HMBC spectrum of siphonellamide B (**2**)

Figure S12. ^1H - ^1H COSY spectrum of siphonellamide B (**2**)

Figure S13. ^1H - ^1H TOCSY spectrum of siphonellamide B (**2**)

Figure S14. HR-ESI-MS spectrum of siphonellamide B (**2**)

Figure S15. ^1H NMR spectrum (500 MHz) of siphonellamide C (**3**) in CDCl_3

Figure S16. HMQC spectrum of siphonellamide C (**3**)

Figure S17. HMBC spectrum of siphonellamide C (**3**)

Figure S18. ^1H - ^1H COSY spectrum of siphonellamide C (**3**)

Figure S19. ^1H – ^1H TOCSY spectrum of siphonellamide C (**3**)

Figure S20. HR-ESI-MS spectrum of siphonellamide C (**3**)

Figure S21. ^1H NMR spectrum (500 MHz) of siphonellamide D (**4**) in CDCl_3

Figure S22. HMQC spectrum of siphonellamide D (**4**)

Figure S23. HMBC spectrum of siphonellamide D (4)

Figure S24. ^1H - ^1H COSY spectrum of siphonellamide D (4)

Figure S25. ^1H – ^1H TOCSY spectrum of siphonellamide D (**4**)

Figure S26. HR-ESI-MS spectrum of siphonellamide D (**4**)

Figure S27. ¹H NMR spectrum (400 MHz) of siphonellamide E (5) in CDCl₃

Figure S28. ¹³C NMR spectrum (100 MHz) of siphonellamide E (5) in CDCl₃

Figure S29. HMBC spectrum of siphonellamide E (5)

Figure S30. HMBC spectrum of siphonellamide E (5)

Figure S31. ^1H – ^1H COSY spectrum of siphonellamide E (**5**)

Figure S32. ^1H – ^1H TOCSY spectrum of siphonellamide E (**5**)

Figure S33. HR-ESI-MS spectrum of siphonellamide E (**5**)

Figure S34. Ozonolysis of siphonellamide E (**5**). **A.** ESI-MS of ozonolysis products. **B.** ESI-MS/MS of m/z 301 [M+H]⁺

Figure S35. ^1H NMR spectrum (500 MHz) of siphonellanol A (**6**) in CDCl_3

Figure S36. ^{13}C NMR spectrum (125 MHz) of siphonellanol A (**6**) in CDCl_3

Figure S37. HMQC spectrum of siphonellanol A (**6**)

Figure S38. HMBC spectrum of siphonellanol A (**6**)

Figure S39. ^1H – ^1H COSY spectrum of siphonellanol A (**6**)

Figure S40. ^1H – ^1H TOCSY spectrum of siphonellanol A (**6**)

Figure S41. HR-ESI-MS spectrum of siphonellanol A (6)

Figure S42. ¹H NMR spectrum (500 MHz) of siphonellanol B (7) in CDCl₃

Figure S43. ^{13}C NMR spectrum (100 MHz) of siphonellanol B (**7**) in CDCl_3

Figure S44. HMQC spectrum of siphonellanol B (**7**)

Figure S45. HMBC spectrum of siphonellanol B (7)

Figure S46. ^1H - ^1H COSY spectrum of siphonellanol B (7)

Figure S47. ^1H - ^1H TOCSY spectrum of siphonellanol B (7)

Figure S48. HR-ESI-MS spectrum of siphonellanol B (7)

Figure S49. ^1H NMR spectrum (400 MHz) of siphonellanol C (**8**) in CDCl_3

Figure S50. ^{13}C NMR spectrum (100 MHz) of siphonellanol C (**8**) in CDCl_3

Figure S51. HMBC spectrum of siphonellanol C (**8**)

Figure S52. HMBC spectrum of siphonellanol C (**8**)

Figure S53. ^1H – ^1H COSY spectrum of siphonellanol C (**8**)

Figure S54. ^1H – ^1H TOCSY spectrum of siphonellanol C (**8**)

Figure S55. HR-ESI-MS spectrum of siphonellanol C (**8**)

Figure S56. ¹H NMR spectrum (500 MHz) of tribromiododiphenyl ether (**13**) in CDCl₃

Figure S57. ¹³C NMR spectrum (125 MHz) of tribromoiododiphenyl ether (**13**) in CDCl₃

Figure S58. HMQC spectrum of tribromoiododiphenyl ether (**13**)

Figure S59. HMBC spectrum of tribromoiododiphenyl ether (**13**)

Figure S60. ^1H - ^1H COSY spectrum of tribromoiododiphenyl ether (**13**)

Figure S61. HR-ESI-MS spectrum of tribromoiododiphenyl ether (**13**)

Figure S62. ^1H NMR spectrum (500 MHz) of gelliodesinic acid (**25**) in CDCl_3

Figure S63. ¹³C NMR spectrum (125 MHz) of gelliodesinic acid (25) in CDCl₃

Figure S64. HMQC spectrum of gelliodesinic acid (25)

Figure S65. HMBC spectrum of gelliodesinic acid (**25**)

Figure S66. ^1H - ^1H COSY spectrum of gelliodesinic acid (**25**)

Figure S67. NOESY spectrum of gelliodesinic acid (**25**)

Figure S68. HR-ESI-MS spectrum of gelliodesinic acid (**25**)

Figure S69. ¹H NMR spectrum (400 MHz) of 3-(propan-2-ylidene)indolin-2-one (26) in CDCl₃

Figure S70. ¹³C NMR spectrum (100 MHz) of 3-(propan-2-ylidene)indolin-2-one (26) in CDCl₃

Figure S71. HMQC spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)

Figure S72. HMBC spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)

Figure S73. ^1H - ^1H COSY spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)

Figure S74. ESI-MS spectrum of 3-(propan-2-ylidene)indolin-2-one (**26**)