

# An efficient synthesis of cycloalkane-1,3-dione-2-spirocyclopropanes from 1,3-cycloalkanediones using (1-aryl-2-bromoethyl)dimethylsulfonium bromides: application to a one-pot synthesis of tetrahydroindol-4(5H)-one

Hisanori Nambu,\* Masahiro Fukumoto, Wataru Hirota, Naoki Ono, Takayuki Yakura\*

Graduate School of Medicine and Pharmaceutical Sciences, University of Toyama, Sugitani, Toyama 930-0194, Japan

## ARTICLE INFO

### Article history:

Received  
Received in revised form  
Accepted  
Available online

### Keywords:


Cyclopropane  
1,3-Cyclohexanediones  
Sulfonium salts  
Indole  
One-pot synthesis

## ABSTRACT

An efficient synthesis of cyclohexane- and cyclopentane-1,3-dione-2-spirocyclopropanes from 1,3-cycloalkanediones using sulfonium salts was achieved. The reaction of 1,3-cycloalkanediones with (1-aryl-2-bromoethyl)dimethylsulfonium bromides and powdered  $K_2CO_3$  in EtOAc provided the corresponding spirocyclopropanes in high yields. Furthermore, a one-pot synthesis of tetrahydroindol-4(5H)-one from 1,3-cyclohexanedione was achieved by using the present protocol and a sequential ring-opening cyclization of spirocyclopropane with a primary amine.

2015 Elsevier Ltd. All rights reserved.


Doubly activated cyclopropanes represent versatile intermediates for the synthesis of a variety of carbo- and heterocyclic compounds.<sup>1</sup> In this context, a ring-opening cyclization reaction of doubly activated cyclopropanes with primary amines is one of the most powerful methods for the construction of pyrrole skeletons.<sup>2</sup> Very recently, we reported the first example of the formation of indole skeletons by employing the ring-opening cyclization of spirocyclopropanes. The reaction of cyclohexane-1,3-dione-2-spirocyclopropanes **3**, derived from 1,3-cyclohexanediones **1**, with primary amines proceeded smoothly at room temperature to give high yields of tetrahydroindol-4(5H)-ones **4**, one of which was easily converted to 4-hydroxyindole **5** (Scheme 1).<sup>3</sup> This procedure provides a useful method for the synthesis of 4-hydroxyindoles. However, there is still a need for improvement in the yields of spirocyclopropanes **3** in Rh(II)-catalyzed cyclopropanation<sup>4-6</sup> with 2-diazo-1,3-cyclohexanediones **2** (27–48% yields). The reaction of alkenes and diazo substrates derived from active methylene compounds in the presence of an Rh(II) catalyst is widely employed for the synthesis of a variety of doubly activated cyclopropanes. In the case of the preparation of spirocyclopropanes, the Rh(II)-catalyzed reaction also produced several cyclopropanes easily but in low yield and was accompanied by a large amount of by-products. For example, the reaction of an excess amount of styrene with 2-diazo-1,3-



**Scheme 1.** Synthesis of 4-hydroxyindole **5** employing a ring-opening cyclization of cyclohexane-1,3-dione-2-spirocyclopropane **3** with amine.

cyclohexanedione (**2a**) using a catalytic amount of  $Rh_2(OAc)_4$  gave spirocyclopropane **3a** in 43% yield and tetrahydrobenzofuran-4(5H)-one **6a**<sup>7</sup> as a by-product in 27% yield (Scheme 1).<sup>6</sup> Furthermore, this protocol is a two-step conversion from 1,3-cyclohexanediones **1** and requires a potentially explosive diazo compound. These drawbacks led us to develop a concise and practical route to spirocyclopropanes **3** from 1,3-cyclohexanediones **1**. Herein, we report an efficient synthesis of 1-arylspiro[2.5]octane-4,8-diones **3** using (1-aryl-2-bromoethyl)dimethylsulfonium bromides **7** (Scheme 2).

\* Corresponding author. Tel.: +81-76-434-7555 or 7556; fax: +81-76-434-5053; e-mail: nambu@pha.u-toyama.ac.jp; yakura@pha.u-toyama.ac.jp




**Scheme 2.** Syntheses of spirocyclopropanes **3** from 1,3-cyclohexanediones **1** and sulfonium salts **7**.

The reaction of active methylene compounds such as malonate and  $\beta$ -ketoester with sulfonium salts has been established<sup>8–10</sup> as an alternative approach to doubly activated cyclopropanes. Recently, Chandrasekaran and Gopinath reported that the reaction of 2,4-pentanedione with (2-bromo-1-phenylethyl)dimethylsulfonium bromide (**7a**) and  $K_2CO_3$  in  $CH_2Cl_2/H_2O$  (1:1) gave the corresponding cyclopropane in 65% yield.<sup>9</sup> Lu and co-workers applied this reaction to a cyclic alkanedione system, in which the reaction was conducted using DBU in DMSO, although the yield was very low.<sup>10</sup> At the outset, we examined the reaction of 1,3-cyclohexanedione (**1a**) with sulfonium salt **7a** for the synthesis of 1-phenylspiro[2.5]octane-4,8-dione (**3a**) according to Chandrasekaran's procedure (Table 1, entry 1). The reaction of **1a** with 1.5 equiv of **7a** using  $K_2CO_3$  in  $CH_2Cl_2/H_2O$  (1:1) at room temperature provided **3a** in 30% yield. To enhance the yield of **3a**, we then screened other solvents. Switching the solvent to  $CH_2Cl_2$  made the reaction messy to give tetrahydrobenzofuran-4(5*H*)-one **6a** in 6% yield without **3a** (entry 2). The use of  $CH_3CN$ , *i*PrOH, and THF resulted in low yields of **3a** (7–14% yields, entries 3–5). We considered that the poor solubility of sulfonium salt **7a** led to low product yields. With the highly polar solvents DMF and DMSO, spirocyclopropane **3a** was obtained in 26% and 35% yields, respectively, along with **6a** in 28% and 23% yields, respectively (entries 6 and 7). Surprisingly, the reaction in EtOAc proceeded smoothly to completion within 1.5 h and gave **3a** in 81% yield (entry 8), although **7a** is almost insoluble in EtOAc. A survey of bases revealed that  $K_2CO_3$  was the optimal base for this transformation (entry 8 vs entries 9 and 10). To our delight, the use of powdered  $K_2CO_3$ , which was finely ground in a mortar, instead of granular  $K_2CO_3$  increased the product yield of **3a** (92% yield, entry 11).

**Table 1**

Reaction of 1,3-cyclohexanedione (**1a**) with sulfonium salt **7a**<sup>a</sup>



entry	base	solvent	time (h)	yield (%) <sup>b</sup>	
				<b>3a</b>	<b>6a</b>
1	$K_2CO_3$	$CH_2Cl_2/H_2O$ (1:1)	8	30	trace
2	$K_2CO_3$	$CH_2Cl_2$	24	trace	6
3	$K_2CO_3$	$CH_3CN$	9	14	trace
4	$K_2CO_3$	<i>i</i> PrOH	20	9	9
5	$K_2CO_3$	THF	24	7	trace
6	$K_2CO_3$	DMF	4	26	28
7	$K_2CO_3$	DMSO	2	35	23
8	$K_2CO_3$	EtOAc	1.5	81	trace
9	<i>t</i> BuOK	EtOAc	24	trace	trace
10	DBU	EtOAc	0.5	20	7
11	$K_2CO_3$ <sup>c</sup>	EtOAc	1	92	trace

<sup>a</sup> All reactions were performed on a 0.5 mmol scale.


<sup>b</sup> Isolated yield.

<sup>c</sup> Powdered  $K_2CO_3$  was used.

With the optimal conditions in hand,<sup>11</sup> we investigated the scope of the reaction of a variety of 1,3-cyclohexanediones **1a–d** with several sulfonium salts **7a–c** prepared from styrene derivatives and bromodimethylsulfonium bromide<sup>12,13</sup> (Table 2). The reaction of **1a** with **7b** and **7c** bearing *p*-bromo or *p*-methyl substituents on the benzene ring provided the corresponding spirocyclopropanes **3b** and **3c**<sup>14</sup> in 92% and 74% yields, respectively. High yields of **3d–f**<sup>14</sup> (84–94% yields) were consistently obtained in the reaction of dimedone (**1b**) with **7a–c**. The reaction of 5-methyl- and 5-phenyl-1,3-cyclohexanediones (**1c** and **1d**) with **7a** afforded diastereomeric mixtures (ca. 1:1) of spirocyclopropanes **3g** and **3h** in 89% and 81% yields, respectively. We then examined the Rh(II)-catalyzed cyclopropanation of the corresponding styrene derivatives with 2-diazo-1,3-cyclohexanediones derived from **1a–d**, and the yields are shown in square brackets in Table 2. In all cases, the yields of **3a–h** (38–47% yields) were lower than those obtained by the reactions of **1** and **7**.

**Table 2**

Reaction of 1,3-cyclohexanediones **1a–d** with sulfonium salts **7a–c**<sup>a</sup>



<sup>a</sup> All reactions were performed on a 0.5 mmol scale with 1.5 equiv of sulfonium salts **7a–c** and 3 equiv of powdered  $K_2CO_3$  in EtOAc.


<sup>b</sup> Yields in square brackets were obtained in the reactions of large amounts of the corresponding styrene derivatives with 2-diazo-1,3-cyclohexanediones derived from **1a–d** in the presence of 1 mol % of  $Rh_2(OAc)_4$ .

<sup>c</sup> Ref. 3.

<sup>d</sup> The products **3g** and **3h** were isolated as a diastereomeric mixture (ca. 1:1).


We next turned our attention to the reaction of 5-membered carbocycles with sulfonium salt **7a** (Scheme 3). The reaction of 1,3-cyclopentanedione (**8**) with 1.5 equiv of **7a** using powdered  $K_2CO_3$  in EtOAc provided the corresponding spirocyclopropane **9** in 83% yield (determined by <sup>1</sup>H NMR analysis).<sup>15,16</sup> The use of 1,3-indanedione (**10**) as an active methylene compound afforded

spirocyclopropane **11**<sup>17</sup> in 85% yield. Thus, the present protocol can be applied to the synthesis of spiro[2.4]heptane compounds.




**Scheme 3.** Reactions of 1,3-cyclopentanedione (**8**) and 1,3-indanedione (**10**) with sulfonium salt **7a**.

To expand the scope of the present protocol, we then examined the reaction with acyclic 1,3-diones (Scheme 4). The reaction of sulfonium salts **7a** and **7b** with 2 equiv of 2,4-pentanedione (**12**) using powdered K<sub>2</sub>CO<sub>3</sub> in EtOAc provided the corresponding doubly activated cyclopropanes **13a**<sup>9</sup> and **13b**<sup>18</sup> in 80% and 88% yields, respectively. The use of 1,3-diphenyl-1,3-propanedione (**14**) as an acyclic 1,3-dione afforded cyclopropane **15** in 85% yield.<sup>19</sup> Since the yields of **13a** and **15** were higher than those reported in literatures,<sup>9,19</sup> these results clearly demonstrate that the combination of powdered K<sub>2</sub>CO<sub>3</sub> as a base and EtOAc as a solvent is also effective for the reaction with acyclic 1,3-diones.



**Scheme 4.** Reactions of sulfonium salts **7a** and **7b** with acyclic 1,3-diones **12** and **14**.

Finally, we investigated a one-pot conversion from 1,3-cyclohexanedione (**1a**) to tetrahydroindol-4(5*H*)-one **4a** (Scheme 5). In our previously reported ring-opening cyclization reaction of spirocyclopropane with a primary amine (Scheme 1), obvious solvent effects were not observed.<sup>3</sup> The reactions were almost the same in CH<sub>2</sub>Cl<sub>2</sub>, CH<sub>3</sub>CN, THF, and toluene. These observations indicate that EtOAc would be usable in this ring-opening cyclization. After the reaction of **1a** with 1.05 equiv of **7a**<sup>20</sup> in EtOAc for 1 h, 2 equiv of benzylamine was directly added to the reaction mixture. The desired ring-opening cyclization reaction proceeded smoothly at room temperature to afford **4a** in 80% yield from **1a**.



**Scheme 5.** One-pot synthesis of tetrahydroindol-4(5*H*)-one **4a** from **1a**.

In conclusion, we have developed an efficient procedure for the synthesis of cyclohexane-1,3-dione-2-spirocyclopropanes from 1,3-cyclohexanediones using sulfonium salts. The reaction of 1,3-cyclohexanediones with (1-aryl-2-bromoethyl)dimethylsulfonium bromides and powdered K<sub>2</sub>CO<sub>3</sub> in EtOAc provided the corresponding spirocyclopropanes in high yields. This method was applicable to 1,3-cyclopentanedione, 1,3-indanedione and acyclic 1,3-diones, affording the corresponding cyclopropanes in high yields. In addition, a one-pot synthesis of tetrahydroindol-4(5*H*)-one from 1,3-cyclohexanedione was achieved by exploiting the present protocol and a sequential ring-opening cyclization of spirocyclopropane with benzylamine. Further application of a one-pot reaction to the synthesis of biologically active indole alkaloids is currently in progress.

## Supplementary Material

Supplementary data associated with this article can be found, in the online version, at <http://...>

## References and notes

- For recent review, see: (a) Carson, C. A.; Kerr, M. A. *Chem. Soc. Rev.* **2009**, *38*, 3051; (b) De Simone, F.; Waser, J. *Synthesis* **2009**, 3353; (c) Campbell, M. J.; Johnson, J. S.; Parsons, A. T.; Pohlhaus, P. D.; Sanders, S. D. *J. Org. Chem.* **2010**, *75*, 6317; (d) Lebold, T. P.; Kerr, M. A. *Pure Appl. Chem.* **2010**, *82*, 1797; (e) Mel'nikov, M. Y.; Budynina, E. M.; Ivanova, O. A.; Trushkov, I. V. *Mendeleev Commun.* **2011**, *21*, 293; (f) Cavitt, M. A.; Phun, L. H.; France, S. *Chem. Soc. Rev.* **2014**, *43*, 804; (g) Schneider, T. F.; Kaschel, J.; Werz, D. B. *Angew. Chem. Int. Ed.* **2014**, *53*, 5504; (h) de Nanteuil, F.; De Simone, F.; Frei, R.; Benfatti, F.; Serrano, E.; Waser, J. *Chem. Commun.* **2014**, *50*, 10912.
- (a) Celerier, J. P.; Haddad, M.; Jacoby, D.; Lhommet, G. *Tetrahedron Lett.* **1987**, *28*, 6597; (b) Jacoby, D.; Celerier, J. P.; Haviari, G.; Petit, H.; Lhommet, G. *Synthesis* **1992**, 884; (c) David, O.; Blot, J.; Bellec, C.; Fargeau-Bellassoued, M.-C.; Haviari, G.; Célérier, J.-P.; Lhommet, G.; Gramain, J.-C.; Gardette, D. *J. Org. Chem.* **1999**, *64*, 3122; (d) Jabin, I.; Monnier-Benoit, N.; Le Gac, S.; Netchitaïlo, P. *Tetrahedron Lett.* **2003**, *44*, 611; (e) Wurz, R. P.; Charette, A. B. *Org. Lett.* **2005**, *7*, 2313; (f) Martin, M. C.; Patil, D. V.; France, S. *J. Org. Chem.* **2014**, *79*, 3030.
- Nambu, H.; Fukumoto, M.; Hirota, W.; Yakura, T. *Org. Lett.* **2014**, *16*, 4012.
- For reviews, see: (a) Lebel, H.; Marcoux, J.-F.; Molinaro, C.; Charette, A. B. *Chem. Rev.* **2003**, *103*, 977; (b) Pellissier, H. *Tetrahedron* **2008**, *64*, 7041; (c) Doyle, M. P. *Angew. Chem. Int. Ed.* **2009**, *48*, 850; (d) Bartoli, G.; Bencivenni, G.; Dalpozzo, R. *Synthesis* **2014**, *46*, 979.
- For selected examples of Rh(II)-catalyzed cyclopropanation, see: (a) Müller, P.; Tohill, S. *Tetrahedron* **2000**, *56*, 1725; (b) Doyle, M. P.; Davies, S. B.; Hu, W. *Org. Lett.* **2000**, *2*, 1145; (c) Yang, M.; Webb, T. R.; Livant, P. *J. Org. Chem.* **2001**, *66*, 4945; (d) Pellicciari, R.; Marinozzi, M.; Camaioni, E.; Nùñez, M. del C.; Costantino, G.; Gasparini, F.; Giorgi, G.; Macchiarulo, A.; Subramanian, N. *J. Org. Chem.* **2002**, *67*, 5497; (e) Doyle, M. P.; Hu, W. *ARKIVOC*, **2003**, *vii*, 15; (f) Wurz, R. P.; Charette, A. B. *Org. Lett.* **2005**, *7*, 2313; (g) Lee, Y. R.; Choi, J. H. *Bull. Korean Chem. Soc.* **2006**, *27*, 503; (h) González-Bobes, F.; Fenster, M. D. B.; Kiau, S.; Kolla, L.; Kolotuchin, S.; Soumeillant, M. *Adv. Synth. Catal.* **2008**, *350*, 813; (i) Marcoux, D.; Charette, A. B. *Angew. Chem. Int. Ed.* **2008**, *47*, 10155; (j) Marcoux, D.; Azzì, S.; Charette, A. B. *J. Am. Chem. Soc.* **2009**, *131*, 6970; (k) Marcoux, D.; Goudreau, S. R.; Charette, A. B. *J. Org. Chem.* **2009**, *74*, 8939; (l) Marcoux, D.; Lindsay, V. N. G.; Charette, A. B. *Chem. Commun.* **2010**, *46*, 910; (m) Lindsay, V. N. G.; Nicolas, C.; Charette, A. B. *J. Am. Chem. Soc.* **2011**, *133*, 8972; (n) Robles, O.; Serna-Saldívar, S.

- O.; Gutiérrez-Urbe, J. A.; Romo, D. *Org. Lett.* **2012**, *14*, 1394; (o) Lindsay, V. N. G.; Fiset, D.; Gritsch, P. J.; Azzi, S.; Charette, A. B. *J. Am. Chem. Soc.* **2013**, *135*, 1463.
6. Müller and co-workers reported that cyclopropanation of styrene with 2-diazodimedone (**2b**) in the presence of Rh<sub>2</sub>(OAc)<sub>4</sub> provided the corresponding spirocyclopropane **3d** in moderate yield (49% yield). Müller, P.; Allenbach, Y. F.; Ferri, M.; Bernardinelli, G. *ARKIVOC*, **2003**, *vii*, 80.
7. (a) Kalpogiannaki, D.; Martini, C.-I.; Nikopoulou, A.; Nyxas, J. A.; Pantazi, V.; Hadjiarapoglou, L. P. *Tetrahedron* **2013**, *69*, 1566; (b) Xia, L.; Lee, Y. R. *Adv. Synth. Catal.* **2013**, *355*, 2361; (c) Bosnidou, A.-E.; Kalpogiannaki, D.; Karanestora, S.; Nixas, J. A.; Hadjiarapoglou, L. P. *J. Org. Chem.* **2015**, *80*, 1279.
8. (a) Gosselck, J.; Béress, L.; Schenk, H. *Angew. Chem. Int. Ed.* **1966**, *5*, 596; (b) Johnson, C. R.; Lockard, J. P. *Tetrahedron Lett.* **1971**, 4589; (c) Johnson, C. R.; Lockard, J. P.; Kennedy, E. R. *J. Org. Chem.* **1980**, *45*, 264; (d) Chow, Y. L.; Bakker, B. H.; Iwai, K. *J. Chem. Soc., Chem. Commun.* **1980**, 521; (e) Jacoby, D.; Celerier, J. P.; Haviari, G.; Petit, H.; Lhommet, G. *Synthesis* **1992**, 884; (f) Kasai, N.; Maeda, R.; Furuno, H.; Hanamoto, T. *Synthesis* **2012**, *44*, 3489; (g) Hirota, K.; Takehiro, Y.; Kamaishi, R.; Yamada, Y.; Hanamoto, T. *Chem. Commun.* **2013**, 49, 7965; (h) Matlock, J. V.; Fritz, S. P.; Harrison, S. A.; Coe, D. M.; McGarrigle, E. M.; Aggarwal, V. K. *J. Org. Chem.* **2014**, *79*, 10226.
9. Gopinath, P.; Chandrasekaran, S. *J. Org. Chem.* **2011**, *76*, 700.
10. Lu and co-workers reported that the reaction of β-(trifluoromethyl)vinyl diphenylsulfonium salt with 1,3-cyclohexanedione (**1a**) in the presence of DBU in DMSO gave 1-trifluoromethylspiro[2.5]octane-4,8-dione in 22% yield. Lin, H.; Shen, Q.; Lu, L. *J. Org. Chem.* **2011**, *76*, 7359.
11. Typical procedure for the synthesis of cyclohexane-1,3-dione-2-spirocyclopropanes **3** with sulfonium salts **7**: 1-Phenylspiro[2.5]octane-4,8-dione (**3a**) (Table 1, entry 11). Powdered K<sub>2</sub>CO<sub>3</sub> (207 mg, 1.5 mmol) and 1,3-cyclohexanedione (**1a**) (56 mg, 0.5 mmol) were added to a suspension of (2-bromo-1-phenylethyl)dimethylsulfonium bromide (**7a**) (245 mg, 0.75 mmol) in EtOAc (5 mL). After stirring at room temperature for 1 h, the reaction mixture was filtered through a Celite pad and the filter cake was rinsed with EtOAc (30 mL). Combined filtrates were washed with water (10 mL) and the aqueous layer was extracted with EtOAc (10 mL x 2). The combined organic layer was washed with brine (10 mL) and dried over anhydrous MgSO<sub>4</sub>. The filtrate was concentrated in vacuo, and the crude product was purified by column chromatography (silica gel, 25% EtOAc in hexane) to provide **3a** (98 mg, 92%) as a white solid.
12. Chow, Y. L.; Bakker, B. H. *Synthesis* **1982**, 648.
13. For reviews, see: (a) Choudhury, M. L. H. *Synlett* **2006**, 1619; (b) Choudhury, L. H.; Parvin, T.; Khan, A. T. *Tetrahedron* **2009**, *65*, 9513.
14. The crude products of **3c** and **3f** were purified by recrystallizations due to the instability of **3c** and **3f** on silica gel.
15. Determined by <sup>1</sup>H NMR analysis of the crude reaction mixture using 1,3,5-trimethoxybenzene as the internal standard.
16. The product **9** was obtained in 54% isolated yield. The crude product of **9** was purified by column chromatography on silica gel despite the instability of **9** on silica gel, because it is very difficult to isolate **9** by recrystallization.
17. Rosenfeld, M. J.; Ravi Shankar, B. K.; Shechter, H. *J. Org. Chem.* **1988**, *53*, 2699.
18. Itoh, O.; Yamamoto, N.; Nakano, K.; Sugita, T.; Ichikawa, K. *Bull. Chem. Soc. Jpn.* **1975**, *48*, 3698.
19. Very recently, Feng and Liu reported that the reaction of sulfonium salts **7a** with 1.2 equiv of 1,3-diphenyl-1,3-propanedione (**14**) using 3 equiv of K<sub>2</sub>CO<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O (1:1) provided the corresponding doubly activated cyclopropane **15** in 68% yield. Xia, Y.; Liu, X.; Zheng, H.; Lin, L.; Feng, X. *Angew. Chem. Int. Ed.* **2015**, *54*, 227.
20. A slightly excess amount (1.05 equiv) of **7a** was used in the one-pot reaction, because sulfonium salts react with primary amines to give aziridines. See ref. 8b–d,g and as follows: (a) Matsuo, J.; Yamanaka, H.; Kawana, A.; Mukaiyama, T. *Chem. Lett.* **2003**, *32*, 392; (b) Yamanaka, H.; Matsuo, J.; Kawana, A.; Mukaiyama, T. *ARKIVOC*, **2004**, *iii*, 42; (c) Maeda, R.; Ooyama, K.; Anno, R.; Shiosaki, M.; Azema, T.; Hanamoto, T. *Org. Lett.* **2010**, *12*, 2548.