

## On the sets of maximum points for generalized Takagi functions

Yasuhiro FUJITA and Yusuke SAITO

**Abstract.** Let  $\varphi$  be a continuous and periodic function on  $\mathbb{R}$  with period 1 and  $\varphi(0) = 0$ . We consider the generalized Takagi function  $f_\varphi$  defined by  $f_\varphi(x) = \sum_{n=0}^{\infty} \frac{1}{2^n} \varphi(2^n x)$  and the set  $M_\varphi$  of maximum points of  $f_\varphi$  in the interval  $[0, 1]$ . When  $\varphi_0(x)$  is the function defined by the distance from  $x$  to the nearest integer,  $f_{\varphi_0}$  is just the Takagi function. Our aim is to seek a condition on  $\varphi$  in order that  $M_\varphi \subset M_{\varphi_0}$ .

### 1. Introduction and the result

The Takagi function is defined by

$$f_{\varphi_0}(x) = \sum_{n=0}^{\infty} \frac{1}{2^n} \varphi_0(2^n x), \quad x \in \mathbb{R}, \quad (1)$$

where  $\varphi_0(x)$  is the function defined by the distance from  $x$  to the nearest integer (cf. [1, 6]). It is a pathological function in the sense that it is everywhere continuous but nowhere differentiable on  $\mathbb{R}$ . Let  $M_{\varphi_0}$  be the set of maximum points in the interval  $[0, 1]$  for the Takagi function  $f_{\varphi_0}$ . By Kahane [4], we have

$$M_{\varphi_0} = \left\{ \sum_{j=1}^{\infty} \frac{a_j}{4^j} \mid a_j \in \{1, 2\} \right\}. \quad (2)$$

---

2010 *Mathematics Subject Classification.* Primary 26A03, 26A27; Secondary 39B22.

*Key words and phrases.* The Takagi function, Set of maximum points, Concavity.

The first author is supported in part by JSPS KAKENHI # 15K04949.

Thus,  $M_{\varphi_0}$  is uncountable. It is important to note that  $\min M_{\varphi_0} = \frac{1}{3}$  and  $\max M_{\varphi_0} = \frac{2}{3}$ .

As a generalization of the Takagi function, we consider the function  $f_\varphi$  of the form

$$f_\varphi(x) = \sum_{n=0}^{\infty} \frac{1}{2^n} \varphi(2^n x), \quad x \in \mathbb{R}, \quad (3)$$

where  $\varphi$  is a continuous and periodic function on  $\mathbb{R}$  with period 1. We call the function of the form (3) a generalized Takagi function. The functions of the form (3) have been considered since the work of [5]. Behrend [3] studied non-differentiability of the functions of the form (3).

In the following, for the function  $f_\varphi$  of (3), we use the notations

$$M_\varphi = \{x \in [0, 1] \mid f_\varphi(x) = m_\varphi\}, \quad m_\varphi = \max_{y \in [0, 1]} f_\varphi(y). \quad (4)$$

Note that  $f_\varphi$  is also periodic with period 1.

Our aim is to compare  $M_\varphi$  with  $M_{\varphi_0}$ . In particular, we are interested in a condition on  $\varphi$  in order that  $M_\varphi \subset M_{\varphi_0}$ .

For a function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ , we consider the following conditions:

- (C1)  $\varphi$  is continuous and periodic on  $\mathbb{R}$  with period 1 and  $\varphi(0) = 0$ . Furthermore,  $\varphi \not\equiv 0$ .
- (C2)  $\varphi(x) = \varphi(1 - x)$ ,  $x \in [0, 1]$ .
- (C3)  $\varphi$  is concave on  $[0, 1]$ , that is, for all  $x, y \in [0, 1]$  and  $\theta \in [0, 1]$ ,

$$\theta\varphi(x) + (1 - \theta)\varphi(y) \leq \varphi(\theta x + (1 - \theta)y).$$

- (C4)  $\varphi(x) < \varphi\left(\frac{1}{3}\right)$ ,  $x \in \left[0, \frac{1}{3}\right)$ .

There exist many functions  $\varphi$  with (C1), (C2), (C3) and (C4).

**Example 1.1.** Let  $\alpha \in \left[\frac{1}{3}, \frac{1}{2}\right]$ . Define the periodic function  $\eta_\alpha : \mathbb{R} \rightarrow [0, \infty)$  with period 1 by  $\eta_\alpha(x) = \min\{\varphi_0(x), \alpha\}$  for  $x \in \mathbb{R}$ . Note that  $\eta_{1/2} \equiv \varphi_0$ . It is easy to see that, for each  $\alpha \in \left[\frac{1}{3}, \frac{1}{2}\right]$ ,  $\eta_\alpha$  fulfills (C1), (C2), (C3) and (C4).

**Theorem 1.2.** *Let  $\varphi$  be a function with (C1), (C2), (C3) and (C4). Then,  $M_\varphi \subset M_{\varphi_0}$ .*

Now, we consider the conditions (C1), (C2), (C3) and (C4). The condition (C1) is natural. The condition (C2) implies that  $\varphi$  is symmetric with respect to the line  $x = \frac{1}{2}$ . The condition (C3) is necessary in Theorem 1.2, since we have an example of a function  $\hat{\varphi}$  with (C1), (C2) and (C4) such that  $\hat{\varphi}$  does not fulfill (C3) and  $M_{\hat{\varphi}} \not\subset M_{\varphi_0}$  (see Example 3.1 below). The condition (C4) is necessary in Theorem 1.2, since we have an example of a function  $\hat{\eta}$  with (C1), (C2) and (C3) such that  $\hat{\eta}$  does not fulfill (C4) and  $M_{\hat{\eta}} \not\subset M_{\varphi_0}$  (see Example 3.2 below).

The contents of the present paper are as follows: In Section 2, we prove Theorem 1.2. In Section 3, we provide two examples which show that the conditions (C3) and (C4) for  $\varphi$  are necessary in order that  $M_\varphi \subset M_{\varphi_0}$ .

## 2. Proof of Theorem 1.2

In this section, we prove Theorem 1.2. We use the notations of Section 1.

**Lemma 2.1.** *Assume that  $\varphi$  fulfills (C1) and (C3). Then,  $\varphi > 0$  in the open interval  $(0, 1)$ .*

*Proof.* We derive a contradiction by supposing that there exists an  $x_0 \in (0, 1)$  such that  $\varphi(x_0) = 0$ . Since  $\varphi \not\equiv 0$  by (C1), we find an  $x_1 \in (0, 1) \setminus \{x_0\}$  such that  $\varphi(x_1) \neq 0$ . First, we consider the case that  $0 < x_1 < x_0$ . By (C3), we have

$$\varphi(x_1) = \varphi\left(\frac{x_1}{x_0}x_0 + \left(1 - \frac{x_1}{x_0}\right) \cdot 0\right) \geq \frac{x_1}{x_0}\varphi(x_0) + \left(1 - \frac{x_1}{x_0}\right)\varphi(0) = 0.$$

Thus,  $\varphi(x_1) > 0$ . On the other hand, let  $\theta = \frac{1-x_0}{1-x_1}$ . Then,  $0 < \theta < 1$  and  $\theta x_1 + (1 - \theta) \cdot 1 = x_0$ . Thus, we have

$$0 = \varphi(x_0) = \varphi(\theta x_1 + (1 - \theta) \cdot 1) \geq \theta\varphi(x_1) + (1 - \theta)\varphi(1) = \theta\varphi(x_1) > 0.$$

This is a contradiction. When  $x_0 < x_1 < 1$ , we can derive a contradiction similarly. Therefore,  $\varphi > 0$  in  $(0, 1)$ .  $\square$

In the following, we assume that  $\varphi$  fulfills (C1), (C2) and (C3), otherwise stated. We define the function  $p_\varphi : \mathbb{R} \rightarrow \mathbb{R}$  of period 1 by

$$p_\varphi(x) = \varphi(x) + \frac{1}{2}\varphi(2x), \quad x \in \mathbb{R}. \quad (5)$$

Note that

$$f_\varphi(x) = \sum_{n=0}^{\infty} \frac{1}{4^n} p_\varphi(4^n x), \quad x \in \mathbb{R}. \quad (6)$$

Let

$$E_\varphi^{(n)} = \{x \in [0, 1] \mid p_\varphi(4^n x) = \mu_\varphi\}, \quad n \in \mathbb{N} \cup \{0\}, \quad (7)$$

where  $\mu_\varphi = \max_{y \in [0, 1]} p_\varphi(y)$ . By (6) and (7), it is clear to see that

$$\bigcap_{n=0}^{\infty} E_\varphi^{(n)} \subset M_\varphi. \quad (8)$$

It is also clear that if  $\bigcap_{n=0}^{\infty} E_\varphi^{(n)} \neq \emptyset$ , the inclusion relation in (8) is reduced to the equality.

Since the following two lemmas are clear, we omit their proofs.

**Lemma 2.2.** *For each  $m \in \mathbb{N}$ , there exist integers  $q_m \in \mathbb{N}$  and  $r_m \in \mathbb{N} \cup \{0\}$  such that*

$$\frac{1}{3}(4^{2m} - 1) = 5q_m, \quad \frac{1}{3}(4^{2m-1} - 1) = 20r_m + 1.$$

**Lemma 2.3.** *Assume that  $\varphi$  fulfills (C1), (C2) and (C3). Then,*

$$f_\varphi(x) = f_\varphi(1-x), \quad p_\varphi(x) = p_\varphi(1-x), \quad x \in [0, 1].$$

**Lemma 2.4.** *Assume that  $\varphi$  fulfills (C1), (C2) and (C3). Then,*

$$p_\varphi(x) \leq p_\varphi\left(\frac{1}{3}\right) = p_\varphi\left(\frac{2}{3}\right), \quad x \in [0, 1].$$

*Proof.* Let  $0 \leq x \leq \frac{1}{2}$ . By (C2) and (C3), we have

$$\begin{aligned} p_\varphi(x) &= \frac{3}{2} \left[ \frac{2}{3}\varphi(1-x) + \frac{1}{3}\varphi(2x) \right] \leq \frac{3}{2} \varphi\left(\frac{2}{3}(1-x) + \frac{1}{3}(2x)\right) \\ &= \frac{3}{2} \varphi\left(\frac{2}{3}\right) = \varphi\left(\frac{2}{3}\right) + \frac{1}{2}\varphi\left(2 \cdot \frac{1}{3}\right) = \varphi\left(\frac{1}{3}\right) + \frac{1}{2}\varphi\left(2 \cdot \frac{1}{3}\right) = p_\varphi\left(\frac{1}{3}\right). \end{aligned}$$

By Lemma 2.3, we conclude the lemma.  $\square$

**Proposition 2.5.** *Assume that  $\varphi$  fulfills (C1), (C2) and (C3). Then,*

$$\frac{1}{3}, \frac{2}{3} \in M_\varphi. \quad (9)$$

*Proof.* By Lemma 2.4, we have  $\frac{1}{3} \in E_\varphi^{(0)}$ . Let  $n \in \mathbb{N}$ . When  $n$  is even, it follows from Lemma 2.2 that there exists an integer  $q_n \in \mathbb{N}$  such that  $4^n \cdot \frac{1}{3} - 5q_n = \frac{1}{3} \in E_\varphi^{(0)}$ . Thus, we have  $\frac{1}{3} \in E_\varphi^{(n)}$ . When  $n$  is odd, it follows from Lemma 2.2 that there exists an integer  $r_n \in \mathbb{N} \cup \{0\}$  such that  $4^n \cdot \frac{1}{3} - (20r_n + 1) = \frac{1}{3} \in E_\varphi^{(0)}$ . Thus, we have  $\frac{1}{3} \in E_\varphi^{(n)}$ . Hence,  $\frac{1}{3} \in E_\varphi^{(n)}$  for all  $n \in \mathbb{N}$ . Therefore,  $\frac{1}{3} \in M_\varphi$  by (8). By Lemma 2.3, we see that  $\frac{2}{3} \in M_\varphi$ .  $\square$

**Proposition 2.6.** *Assume that  $\varphi$  fulfills (C1), (C2) and (C3). Then,*

$M_\varphi = \bigcap_{n=0}^{\infty} E_\varphi^{(n)}$ . Furthermore,  $m_\varphi = 2\varphi\left(\frac{1}{3}\right)$ , where  $m_\varphi$  is the constant of (4).

*Proof.* Note that  $\frac{1}{3} \in \bigcap_{n=0}^{\infty} E_\varphi^{(n)}$  by the proof of Proposition 2.5. Since  $\bigcap_{n=0}^{\infty} E_\varphi^{(n)} \neq \emptyset$ , it is easy to see that  $M_\varphi = \bigcap_{n=0}^{\infty} E_\varphi^{(n)}$  by (8). Furthermore, by (6), (8) and Proposition 2.5, we have

$$m_\varphi = \sum_{n=0}^{\infty} \frac{1}{4^n} p_\varphi\left(\frac{1}{3}\right) = 2\varphi\left(\frac{1}{3}\right).$$

Here, we used  $p_\varphi\left(\frac{1}{3}\right) = \frac{3}{2}\varphi\left(\frac{1}{3}\right)$ .  $\square$

**Proposition 2.7.** *Assume that  $\varphi$  fulfills (C1), (C2), (C3) and (C4).*

*Then,  $\min M_\varphi = \frac{1}{3}$  and  $\max M_\varphi = \frac{2}{3}$ .*

*Proof.* Let  $0 \leq x < \frac{1}{4}$ . Then, by (C3) and (C4), we have

$$\begin{aligned} p_\varphi(x) &= \frac{3}{2} \left[ \frac{2}{3}\varphi(x) + \frac{1}{3}\varphi(2x) \right] \leq \frac{3}{2} \varphi\left(\frac{2}{3}x + \frac{1}{3}(2x)\right) = \frac{3}{2} \varphi\left(\frac{4}{3}x\right) \\ &< \frac{3}{2} \varphi\left(\frac{1}{3}\right) = \varphi\left(\frac{1}{3}\right) + \frac{1}{2} \varphi\left(2 \cdot \frac{1}{3}\right) = p_\varphi\left(\frac{1}{3}\right). \end{aligned}$$

Thus,  $[0, \frac{1}{4}) \cap E_\varphi^{(0)} = \emptyset$ . By Proposition 2.6,  $[0, \frac{1}{4}) \cap M_\varphi = \emptyset$ . For each  $x \in [0, \frac{1}{4})$  and  $n \in \mathbb{N}$ , we have

$$p_\varphi\left(4^n \left(\sum_{i=1}^n \frac{1}{4^i} + \frac{x}{4^n}\right)\right) = p_\varphi(x) < p_\varphi\left(\frac{1}{3}\right).$$

Thus,  $\left[\sum_{i=1}^n \frac{1}{4^i}, \sum_{i=1}^{n+1} \frac{1}{4^i}\right) \cap E_\varphi^{(n)} = \emptyset$ . By Proposition 2.6,  $\left[\sum_{i=1}^n \frac{1}{4^i}, \sum_{i=1}^{n+1} \frac{1}{4^i}\right) \cap M_\varphi = \emptyset$ . Since  $\sum_{i=1}^{\infty} \frac{1}{4^i} = \frac{1}{3}$ , we conclude that  $[0, \frac{1}{3}) \cap M_\varphi = \emptyset$ . Since  $\frac{1}{3} \in M_\varphi$  by Proposition 2.5, we have shown that  $\min M_\varphi = \frac{1}{3}$ . By Lemma 2.3, we have  $\max M_\varphi = \frac{2}{3}$ .  $\square$

**Lemma 2.8.** *Assume that  $\varphi$  and  $\psi$  fulfill (C1), (C2) and (C3). If  $E_\varphi^{(0)} \subset E_\psi^{(0)}$ , then  $M_\varphi \subset M_\psi$ .*

*Proof.* Let  $x \in M_\varphi$ . Then, by Proposition 2.6,  $x \in E_\varphi^{(n)}$  for all  $n \in \mathbb{N} \cup \{0\}$ . Thus,  $p_\varphi(4^n x) = p_\varphi(\frac{1}{3})$  for all  $n \in \mathbb{N} \cup \{0\}$ , which implies that  $4^n x - [4^n x] \in E_\varphi^{(0)}$  for all  $n \in \mathbb{N} \cup \{0\}$ , where  $[y]$  denotes the greatest integer not exceeding  $y \in \mathbb{R}$ . Since  $E_\varphi^{(0)} \subset E_\psi^{(0)}$ , we have  $4^n x - [4^n x] \in E_\psi^{(0)}$  for all  $n \in \mathbb{N} \cup \{0\}$ . Then,  $x \in E_\psi^{(n)}$  for all  $n \in \mathbb{N} \cup \{0\}$ . Hence,  $x \in M_\psi$  by Proposition 2.6.  $\square$

Now, we prove Theorem 1.2.

*Proof of Theorem 1.2.* Let  $\varphi$  be a function with (C1), (C2), (C3) and (C4). By the proof of Proposition 2.7, we have  $[0, \frac{1}{4}) \cap M_\varphi = \emptyset$ , so that  $E_\varphi^{(0)} \subset [\frac{1}{4}, \frac{3}{4}]$  by Lemma 2.3. Since  $E_{\varphi_0}^{(0)} = [\frac{1}{4}, \frac{3}{4}]$ , we conclude the theorem by Lemma 2.8.  $\square$

### 3. Examples

In this section, we provide two examples which show that the conditions (C3) and (C4) for  $\varphi$  are indispensable in order that  $M_\varphi \subset M_{\varphi_0}$ .

**Example 3.1.** Let  $\hat{\varphi} : \mathbb{R} \rightarrow \mathbb{R}$  be the periodic function with period 1 such that  $\hat{\varphi}(x) = (\varphi_0(x))^2$  in  $\mathbb{R}$ . In this case,  $\hat{\varphi}$  fulfills (C1), (C2) and (C4). However,  $\hat{\varphi}$  does not fulfill (C3). We show that  $M_{\hat{\varphi}} \not\subset M_{\varphi_0}$ .

Indeed, we see that

$$\sum_{n=0}^{\infty} \frac{1}{2^n} (\varphi_0(2^n x))^2 = x - x^2, \quad x \in [0, 1],$$

since  $\psi^0(x) = x$  and  $\psi^n(x) = 2\varphi_0(2^{n-1}x)$  ( $n \in \mathbb{N}$ ) in the notations of the example of [8, p.335]. Thus,  $f_{\hat{\varphi}}(x) = x - x^2$  in  $[0, 1]$ . Hence,  $M_{\hat{\varphi}} = \{\frac{1}{2}\}$ .

**Example 3.2.** Define the periodic function  $\hat{\eta} : \mathbb{R} \rightarrow [0, \infty)$  with period 1 by  $\hat{\eta}(x) = \min\{\varphi_0(x), \frac{1}{15}\}$  for  $x \in \mathbb{R}$ . In this case,  $\hat{\eta}$  fulfills (C1), (C2) and (C3). However,  $\hat{\eta}$  does not fulfill (C4). We show that  $M_{\hat{\eta}} \not\subset M_{\varphi_0}$ .

Indeed, we have, on  $[0, \frac{1}{2}]$ ,

$$p_{\hat{\eta}}(x) = \begin{cases} 2x, & 0 \leq x \leq \frac{1}{30}, \\ x + \frac{1}{30}, & \frac{1}{30} \leq x \leq \frac{1}{15}, \\ \frac{1}{10}, & \frac{1}{15} \leq x \leq \frac{7}{15}, \\ -x + \frac{17}{30}, & \frac{7}{15} \leq x \leq \frac{1}{2}. \end{cases}$$

Thus,  $E_{\hat{\eta}}^{(0)} = [\frac{1}{15}, \frac{7}{15}] \cup [\frac{8}{15}, \frac{14}{15}]$ .

We show that  $\frac{1}{15} \in M_{\hat{\eta}}$ . Note that  $\frac{1}{15} \in E_{\hat{\eta}}^{(0)}$ . Fix  $n \in \mathbb{N}$  arbitrarily. When  $n$  is even, it follows from Lemma 2.2 that there exists a number  $q_n \in \mathbb{N}$  such that  $4^n \cdot \frac{1}{15} = q_n + \frac{1}{15}$ . Since  $\frac{1}{15} \in E_{\hat{\eta}}^{(0)}$ , we have  $\frac{1}{15} \in E_{\hat{\eta}}^{(n)}$ . When  $n$  is odd, it follows from Lemma 2.2 that there exists a number  $r_n \in \mathbb{N} \cup \{0\}$  such that  $4^n \cdot \frac{1}{15} = 4r_n + \frac{4}{15}$ . Since  $\frac{4}{15} \in E_{\hat{\eta}}^{(0)}$ , we have  $\frac{1}{15} \in E_{\hat{\eta}}^{(n)}$ . Therefore,  $\frac{1}{15} \in E_{\hat{\eta}}^{(n)}$  for all  $n \in \mathbb{N}$ , and  $\frac{1}{15} \in M_{\hat{\eta}}$ . Since  $\min M_{\varphi_0} = \frac{1}{3}$ , we see that  $\frac{1}{15} \notin M_{\varphi_0}$ . Thus, we conclude the assertion.

### References

- [1] P. ALLAART AND K. KAWAMURA, *The Takagi function: a survey*, Real Anal. Exchange 37 (2011/12), 1–54.

- [2] Y. BABA , *On maxima of Takagi-van der Waerden functions* , Proc. Amer. Math. Soc. 91 (1984), 373–376.
- [3] F. A. BEHREND , *Some remarks on the construction of continuous non-differentiable functions* , Proc. London Math. Soc. 50, (1949), 463–481.
- [4] J.-P. KAHANE, *Sur l'exemple, donné par M. de Rham, d'une fonction continue sans dérivée*, Enseignement Math., 5 (1959), 53–57.
- [5] K. KNOPP, *Ein einfaches Verfahren zur Bildung stetiger nirgends differenzierbarer Funktionen*, Math. Z. 2 (1918), 1–26.
- [6] J. C. LAGARIAS, *The Takagi function and its properties*, RIMS Kôkyûroku Bessatsu, B34 (2012), 153–189.
- [7] T. TAKAGI, *A simple example of the continuous function without derivative*, Phys.-Math. Soc. Japan 1 (1903), 176–177. The Collected Papers of Teiji Takagi, S. Kuroda, Ed., Iwanami (1973), 5–6.
- [8] M. YAMAGUTI AND M. HATA, *Weierstrass's function and chaos*, Hokkaido Math. J. 12 (1983), 333–342.

Yasuhiro Fujita  
Department of Mathematics  
Faculty of Science  
University of Toyama  
Gofuku, Toyama 930-8555, JAPAN  
e-mail: yfujita@sci.u-toyama.ac.jp

Yusuke Saito  
Graduate School of Science and Engineering for Education  
University of Toyama  
Gofuku, Toyama 930-8555, JAPAN  
e-mail: d1771001@ems.u-toyama.ac.jp

(Received July 12, 2017)