

学位論文

**Changes in the association between postpartum depression and
mother-infant bonding by parity: Longitudinal results from the
Japan Environment Children's Study**

産後うつとボンディングの関連の経産による変化
子どもの健康と環境に関する全国調査からの経時的な結果より

富山大学学術研究部医学系 公衆衛生学講座 助教

土田暁子

Changes in the association between postpartum depression and mother-infant bonding by parity: longitudinal results from the Japan Environment and Children's Study

Akiko Tsuchida, MSc

Department of Public Health, Faculty of Medicine, University of Toyama, Toyama, Japan

²Toyama Regional Center for JECS, University of Toyama, Toyama, Japan

Abstract

The results of several epidemiological studies have shown a moderate association between postpartum depression (PPD) and mother-infant bonding (MIB); however, associations and changes that longitudinally considered parity were not reported. We investigated the possible association between PPD and MIB at one month after birth, in addition, the changes of these indices by parity. From a dataset comprising 103,099 maternal registrations in The Japan Environment and Children's Study, the present study analyzed complete data on questionnaires for 76,363 women who participated once (cross-sectional group) and 3,753 women who participated twice (follow-up subgroup). Edinburgh Postnatal Depression Scale (EPDS) scores and the Mother-to-Infant Bonding Scale-Japan (MIBS-J) scores were obtained one month after delivery. We identified the

two factors of the MIBS-J, "lack of maternal feeling (LMF)" and "anxiety about caregiving (AC)" through confirmatory factor analysis. Associations between total EPDS and each factor of MIBS-J were evaluated using multiple regression analyses after adjusting for potential confounders. Total EPDS and both factors (LMF and AC) were positively related in the cross-sectional group (fully adjusted $\beta = 0.26$ and 0.39 for LMF and AC, respectively), in the follow-up subgroup at first participation (0.24 and 0.40 , respectively) and at the second participation (0.25 and 0.39 , respectively). Multiple regression analyses revealed a robust, moderate relationship between postpartum depression and mother-infant bonding. PPD and MIB scores were shown to decrease from the first child to the second in the follow-up subgroup. Consequently, interventions which would increase a mother's child care experience and expertise would prevent PPD and improve MIB.

Keywords: Postpartum depression; Bonding; Maternal attachment; Parity; Parenting; Birth cohort

Abbreviations

PPD: Postpartum Depression; MIB: Mother-Infant Bonding; EPDS: Edinburgh Postnatal Depression Scale; MIBS-J: Mother-to-Infant Bonding Scale-Japan; LMF: Lack of Maternal Feeling; AC: Anxiety about Caregiving; JECS: The Japan Environment and Children's Study; EFA: Exploratory Factor Analysis; CFA: Confirmatory Factor Analysis; CFI: Comparative Fit Index; RMSEA: Root Mean Square Error of Approximation; MI: Multiple Imputation

Introduction

Immediately after giving birth, most mothers experience a special surge of affection for their infants and automatically begin nursing them and keeping them clean. However, in some cases, the mother does not experience these feelings, or she may even feel an aversion to the child. Consequently, she is unable to bond with the child or provide it with adequate maternal care. Suggested risk factors include being unmarried or unemployed (Figueiredo et al., 2009), the spousal relationship (van Bussel et al., 2010), having a Caesarean section (Sockol et al., 2014) the trait anger of mother's personality (Kitamura et al., 2013), social support during pregnancy (Ohara et al., 2017; Ohara et al., 2018), and a mother's perceptions of how she was reared as a child (Ohara et al., 2018). In addition, suggested child-related risk factors include crying at night (Yalcin et al., 2010) and, in cultures with a preference for boys, being a girl (Edhborg et al., 2011).

Much of this research has indicated an association between postpartum depression (PPD) and mother-infant bonding (MIB) difficulties (Dubber et al., 2015; Lefkovic et al., 2014; Muzik et al., 2013; O'Higgins et al., 2013; Ohoka et al., 2014; Sockol et al., 2014). Given that psychotherapy has been reported to be effective in treating PPD (Cuijpers et al., 2008; Dennis and Hodnett, 2007), such treatment may alleviate or prevent PPD and associated MIB difficulties. On the other hand, not all bonding difficulties are related to PPD, making it vital to better understand that relationship and other factors that may be involved.

Studies in Japan have shown that primipara women are at a significantly higher risk of PPD than multiparous women (Mori et al., 2016; Morikawa et al., 2015; Takehara et al., 2018) suggesting that, as the number of births increases, the frequency of PPD and its associated MIB difficulties may decrease. However, no studies have examined the effect of increasing parity on both PPD and MIB difficulties across births in the same mothers.

The Japan Environment and Children's Study (JECS) was a nationwide birth cohort study where 103,099 pregnant women registered to participate between 2011–2014. The first part of the present study took a cross-sectional look at the relationship between PPD and MIB in responses collected from these participants one month after giving birth. During the JECS's 2011–2014 recruitment period, it was possible for participants who

became newly pregnant to re-register to participate again. Consequently, 5,597 (5.4%) of the women participated in the study twice, and 48 (0.05%) participated three times. Taking the opportunity to do an unprecedented study on the effect of parity in the same mothers, the second part of the present study examined the relationship between PPD and MIB. In addition, the changes of these indices as parity increased was evaluated by comparing the first and second participation for women who gave birth twice during the study.

This thesis is based on the published article (Tsuchida et al., 2019).

Methods

Study design

The JECS is a birth cohort study, which investigated the links between environmental factors and childhood health. Recruitment for the study occurred across fifteen regional centers within Japan from 2011 to 2014. Participant recruitment involved a face-to-face explanation of the survey to mothers, and self-administered informed consent was obtained. The details of the JECS design are well documented (Kawamoto et al., 2014; Michikawa et al., 2018).

The JECS protocol received approval from the Ethics Committees of all participating institutions and the Ministry of the Environment.

Study data

The present study used the dataset jecs-ag-20160424, which was released in June 2016 and revised in October 2016, along with the supplementary dataset jecs-ag-20160424-sp1. To arrive at the number of unique mothers who had a singleton live birth, from the data for the 103,099 registrants, we eliminated the data for the second and third births of the 5,645 who had participated more than once, the data from 29 participants who withdrew from the study, the data from 949 participants who had multiple births, and the data from 3,680 participants whose pregnancies ended in miscarriages or still births. Consequently, data from 92,796 mothers remained. After eliminating cases with insufficiently complete responses, the data from 76,373 mothers were analyzed as the “cross-sectional group” for the first analysis. Further, women who participated twice or more ($n = 3,753$) formed the “follow-up subgroup” (Fig. 1).




Figure 1. Participant flow diagram

Measurements

A self-administered questionnaire was administered to the mothers on 3 occasions—the first trimester, the second/third trimester, and one month after delivery—to collect demographics, medical and obstetric history, physical and mental health, lifestyle, occupation, and socioeconomic status. Questionnaires at one month after delivery were distributed by our staff to participants, then collected at the maternity hospital by our staff when participants visited for their one-month-old's health check-up. If the questionnaires could not be collected at that time, they were returned by post.

Questionnaires at one month after delivery included the Edinburgh Postnatal Depression Scale (EPDS) Japanese version (Okano et al., 1996) and 5 items from the Japanese version of the Mother-to-Infant Bonding Scale (MIBS-J) (Yoshida et al., 2012).

The EPDS consisted of 10 items, which were answered using a four-point scale (range = 0–30). We regarded a total EPDS score ≥ 9 as being indicative of PPD. This cut-off score of nine has been validated as the optimal threshold for the Japanese population when screened for minor and major depressive episodes among postpartum women, and has been found to have a sensitivity of 75% and 82% for minor and major depressive episodes and a specificity of 93% and 95%, respectively (Okano et al., 1996; Yamashita et al., 2000). Further, a score ≥ 9 has previously been used to indicate PPD in various

prospective studies in Japan (Ishikawa et al., 2011; Kubota et al., 2014; Takehara et al., 2018). Previous research has shown that the EPDS has a three-factor structure that includes anxiety, depression, and anhedonia among Japanese mothers (Kubota et al., 2014). We referred to the factors and used the sum of the relevant items to represent “anxiety” (EPDS item 3, 4, and 5), “depression” (Item 7, 8, and 9), and “anhedonia” (Item 1 and 2) symptoms.

We used five items from the MIBS-J to assess mother-infant bonding. The five items of MIBS-J that have been commonly used by Japanese public health nurses to assess mother-infant bonding (Hirokawa et al., 2018) include the following: Item 1, “I feel loving towards my baby”(reversed scored); Item 2, “I feel scared or panicky when I have to do something for my baby”; “Item 3, “I feel nothing for my baby”; Item 4, “I enjoy doing things with my baby” (reversed scored); and Item 5, “I wish I did not have my baby.” Item 2, Item 3 and Item 5 were rated on a four-point scale (0, not at all; 1, slightly, some of the time; 2, very much so, some of the time; and 3, very much so, most of the time). The scores were reversed for Item 1 and Item 4.

We examined the factor structure of the five items of MIBS-J by an exploratory factor analysis (EFA). The factor structure derived from the EFA data was confirmed by a confirmatory factor analysis (CFA). Our two factors structure (the $\chi^2/df= 5.3$, CFI = .954,

and RMSEA = 0.077) had a similar fit to the two factors structure that was previously reported by Yoshida et al. (the $\chi^2/df=6.1$, CFI = .912, and RMSEA = 0.077) (Yoshida et al., 2012). The factor analysis indicated a two factors model consisting of three items (Item 1, 3 and 5), and two items (Item 2 and 4). We named the former factor "lack of maternal feeling (LMF)" and the latter factor "anxiety about caregiving (AC)". Each factor was estimated by summing of their item scores (Fig. 2).




Figure 2. Confirmatory factor analysis of the MIBS-J at one month after birth

Statistical analysis

Data were expressed as means \pm standard deviation or medians unless stated otherwise.

Multiple regression analyses were performed to examine the association between the factors from MIBS-J and total EPDS scores, adjusting for parity in Model 1, also adjusting for parity, mother age, highest educational level, annual household income, marital status, alcohol intake, smoking status, physical activity, history of anxiety disorder, history of anxiety, history of depression, history of schizophrenia, diagnostic record of mental disorder during pregnancy, infants' anomaly, infants' sex, and intensity and frequency of infants' crying in Model 2. Covariates for the model were mentioned below.

Maternal age was treated as a continuous variable. Number of parities was categorized as primipara or multipara. Educational background (highest level of education attained) was initially classified into 7 groups—1, junior high school; 2, high school; 3, technical junior college; 4, technical/vocational college; 5, associate degree; 6, bachelor's degree; and 7, postgraduate degree—and then re-categorized into 3 groups (1 + 2, < 13 y; 3 + 4 + 5, 13-14 y; or 6 + 7, > 14y). Annual family income was originally classified into 9 groups and then re-categorized into 3 groups (< 4 million, 4–6 million, or \geq 6 million Japanese yen). Marital status was initially classified into 4 groups—1, married (including

common-law status); 2, single (never married); 3, divorced; and 4, widowed)—and then re-categorized into 3 groups (1, Married; 2, Single; or 3 + 4, Divorced or widowed). Alcohol intake was classified into 6 groups for 1 month after delivery (1, do not drink; 2, drank in the past but stopped drinking; 3, drinks 1–3 times/month; 4, drinks 1–3 times/week; 5, drinks 4–6 times/week; and 6, drinks every day) and then re-categorized into 3 groups (1, Never drinker; 2, Ex drinker; or 3 + 4 + 5 + 6, Current drinker). Smoking status was classified into 6 groups at 1 month after delivery (1, never; 2, smoked previously but quit before learning of current pregnancy; 3, smoked previously, but quit after learning of current pregnancy; 4, still smoking 1–10 cigarettes/day; 5, still smoking 11–20 cigarettes/day; and 6, still smoking ≥ 21 cigarettes/day) and then re-categorized into 3 groups (1, Never smoker; 2 + 3, Ex-smoker; or 4 + 5 + 6, Current smoker). There were no questions on physical activity during the postpartum period; therefore, data from the second/third trimester were used. Physical activity was defined as walking for at least 10 min continuously per week (*yes* or *no*). History of anxiety, depression, schizophrenia, and diagnostic record of mental disorder during pregnancy were categorized as *yes* or *no*. Infant’s anomaly was categorized as *yes* or *no*, that congenital anomalies from the supplementary dataset jecs-ag-20160424-sp1, were included or not. Infants’ sex was

categorized as male or female. Intensity and frequency of infants' crying was classified into 3 groups—1, quite often and long; 2, sometimes and short; and 3, hardly.

Little's Missing Completely at Random test was found to be significant ($p < .001$) for the cross-sectional data set before exclusion of cases due to incomplete responses ($N = 92,796$), supporting the theory that the data was considered missing not at random (Little, 1988). Multiple imputation (MI) was used to account for incomplete answers. The 18 variables included in Model 2 of multiple regression analysis were used in imputation based on chained equations (van Buuren, 2007); 200 imputed data sets were constructed. We compared the standardized beta from multiple regression models applied to the imputed data ($N = 92,796$) and the data of complete cases across the all covariates ($N = 76,373$).

Statistically significant differences were determined by Welch's t -test or paired t -tests to compare means. Two-tailed p -values $< .05$ were considered statistically significant. Data were analyzed using SAS version 9.4 software (SAS Institute Inc., Cary, NC) and IBM SPSS Statistics, version 25 (IBM, Armonk, NY).

Results

Prevalence of PPD and characteristics of participants

Table 1 shows demographic and obstetric characteristics for the cross-sectional group (N = 76,373), and the first and second participations of the follow-up subgroup (n = 3,753). In the cross-sectional group, 10,551 (13.8%) mothers had an EPDS score ≥ 9 . Further, analysis of the 3,753 participants in the follow-up subgroup showed that 541 (14.4%) and 383 (10.2%) had scores ≥ 9 after the first and second participation, respectively (Supplementary Table 1).

The mean of maternal age and the rate of married about the follow-up subgroup increased from the first to the second participation, as expected. Moreover, the highest educational level, alcohol intake, smoking status, history of anxiety disorder, history of depression and history of anxiety disorder were almost same rate of the first and the second participation.

Table 1 Demographic and obstetric characteristics of participants

Characteristics	Cross-sectional group (N = 76,373)	
	Mean / n	(SD / %)
Maternal age, mean (SD)	31.5	(4.9)
Postpartum depression, n (%)	10,551	(13.8)
Previous deliveries, n (%)		
Nullipara	32,342	(42.4)
Multipara	44,031	(57.7)
Highest educational level (year), n (%)		
<13	26,587	(34.8)
13-14	32,524	(42.6)
≥15	17,262	(22.6)
Annual household income (JPY), n (%)		
<4 million yen	30,288	(39.7)
4-5.9 million yen	25,425	(33.3)
≥6 million yen	20,660	(27.1)
Marital status, n (%)		
Married (including common law marriage)	73,423	(96.1)
Single (never married)	2,337	(3.1)
Divorced or widowed	613	(0.8)
Alcohol intake, n (%)		
Never drinker	69,707	(91.3)
Ex drinker	3,450	(4.5)
Current drinker	3,216	(4.2)
Smoking status, n (%)		
Never smoker	44,985	(58.9)
Ex smoker	28,286	(37.0)
Current smoker	3,102	(4.1)
Physical activity, yes (%)	54,974	(72.0)
History of anxiety disorder, yes (%)	2,144	(2.8)
History of depression, yes (%)	2,258	(3.0)
History of schizophrenia, yes (%)	127	(0.2)
Diagnostic record of mental disorder during pregnancy, yes (%)	589	(0.8)
Infant's anomaly, yes (%)	1,522	(2.0)
Infant's gender, n (%)		
Male	39,106	(51.2)
Female	37,267	(48.8)
Intensity and frequency of infant's crying, n (%)		
Quite often and long	12,844	(16.8)
Sometimes and short	60,000	(78.6)
Hardly	3,529	(4.6)

Multivariable regression for PPD and MIB

Table 2 shows the results of multiple regression analyses performed to look at the relationship between total EPDS scores and LMF or AC scores for the cross-sectional group. Total EPDS scores and factor scores were positively related in the unadjusted model and the adjusted models. In the complete case analysis based on 76,373 cases, the relationship between total EPDS and LMF scores were almost unchanged among the unadjusted model, Model 1, and Model 2 ($\beta = 0.27, 0.27$ and 0.26 , respectively). On the other hand, the standardized-betas from the regression model between total EPDS and AC scores were changed by adjusted parity ($\beta = 0.43$ and 0.39 , respectively). However, there was no change by adjusted parity or fully adjusted ($\beta = 0.39$ and 0.39 , respectively).

In the multiple imputation analysis based on 92,769 cases, the relationship between total EPDS and LMF scores are almost unchanged among the unadjusted model, Model 1 and Model 2 ($\beta = 0.29, 0.29$ and 0.28 , respectively). On the other hand, the standardized betas from the regression model between total EPDS and AC scores are the same as the complete case analysis. For follow-up subgroup scores after the first and second participations, total EPDS and the factor's scores were positively related. Further, the fully adjusted standardized beta values for the first and second participation scores were virtually unchanged.

Table 2. Results of multiple regression analyses between postpartum depression (total EPDS score) and bonding factors from MIBS-J at one-month after birth for cross-sectional group

	Cross-sectional group				Follow-up subgroup				
	Complete case analysis (N = 76,373)		Multiple imputation analysis (N = 92,796)		First participation (n = 3,753)		Second participation (n = 3,753)		
	Standardized- beta	<i>p</i>	Standardized- beta	<i>p</i>	Standardized- beta	<i>p</i>	Standardized- beta	<i>p</i>	
Total MIBS-J									
Crude	0.45	<.0001	0.45	<.0001	0.44	<.0001	0.40	<.0001	
Model1	0.41	<.0001	0.42	<.0001	0.42	<.0001	0.40	<.0001	
Model2	0.41	<.0001	0.41	<.0001	0.41	<.0001	0.41	<.0001	
Lack of maternal feeling									
Crude	0.27	<.0001	0.29	<.0001	0.25	<.0001	0.25	<.0001	
Model1	0.27	<.0001	0.29	<.0001	0.24	<.0001	0.25	<.0001	
Model2	0.26	<.0001	0.28	<.0001	0.24	<.0001	0.25	<.0001	
Anxiety about caregiving									
Crude	0.43	<.0001	0.43	<.0001	0.43	<.0001	0.39	<.0001	
Model1	0.39	<.0001	0.39	<.0001	0.41	<.0001	0.39	<.0001	
Model2	0.39	<.0001	0.39	<.0001	0.40	<.0001	0.39	<.0001	

Model 1: adjusted for parity.

Model 2: adjusted for parity, mother's age, highest educational level, annual household income, marital status, alcohol intake, smoking status, physical activity, history of anxiety disorder, history of anxiety, history of depression, history of schizophrenia, diagnostic record of mental disorder during pregnancy, infant's anomaly, infant's sex, intensity and frequency of infant's crying

The difference between EPDS score and MIBS-J scores between the first and second participation of the follow-up subgroup

Table 3 shows the means and standard deviations for the total EPDS scores, the three factors scores from the EPDS (e.g., anxiety, depression and anhedonia), the LMF scores, and the AC scores for the first and second participation scores in the follow-up subgroup. The total EPDS scores, anxiety scores, depression scores, LMF scores, and AC scores were lower in mothers at the time of their second participation than at their first participation.

Table 3. Means (SD) of EPDS score and bonding score in the follow-up subgroup

Name of the scores or questions	Follow-up subgroup (n = 3,753)				
	First participation		Second participation		* <i>P</i>
	mean	(SD)	mean	(SD)	
Total EPDS	5.10	3.51	4.31	3.42	<.0001
Anxiety	2.95	1.96	2.37	1.90	<.0001
Anheadnia	0.17	0.55	0.17	0.57	0.94
Depression	0.65	1.07	0.52	1.04	<.0001
Total MIBS-J	1.51	1.37	0.97	1.22	<.0001
Lack of maternal feeling	0.20	0.60	0.18	0.60	0.04
Anxiety about caregiving	1.31	1.05	0.79	0.88	<.0001

**P* from paired T-test

Discussion

Prevalence of PPD and characteristics of participants

The number of women in the cross-sectional group who had depressive tendencies was consistent with the occurrence found in a systematic review of studies performed in Western countries (around 11% at approximately 1–3 months postpartum; Carroll, Daly, & Begley, 2016) and it fell within the range of 10–15% found in Japanese studies (Choi et al., 2010; Kitamura et al., 2006; Kokubu et al., 2012).

In the follow-up subgroup, some maternal characteristics like the highest educational level, alcohol intake, smoking status and their history of illness are theoretically not changed from the first to the second participation because they are the same person's characteristics. There were small differences between the first and the second participation in these characteristics, however they were not statistically differences. This showed almost all participants in the follow-up subgroup appropriately answered.

The relationship between PPD and MIB

The relationship was calculated between the total EPDS, "lack of maternal feeling" (LMF), and "anxiety about caregiving" (AC) discovered by factor analysis based on the five items from MIBS-J. The results suggest that the PPD (EPDS scores) and bonding

scores have a moderate relationship with Japanese multiparous women. Furthermore, it was revealed that robust results were shown even when adjusted for various modulators without changing the relationship (Table 2).

When conducting the complete case analysis, 20,473 cases were excluded out of 92,796; 92,796 were examined using MI. Compared with the complete case analysis the standardized-beta increased for LMF by approximately 0.02. However, there was no change in the relationship to AC, which suggests that excluding 20,473 cases did not substantially influence the results. Regarding AC, if adjusted by parity, the standardized-beta decreased by 0.04 points, and there is no variation in the fully adjusted scores. Therefore, AC appears to be slightly influenced by parity, but a moderate relationship between PPD and AC remains. As described above, the relationship of PPD and the two factors of MIBS-J showed that the relationship is at the moderate level, which is consistent with prior studies where the relationship was examined one month after birth (Ohoka et al., 2014; Taylor et al., 2005).

A multiple linear regression analysis was also conducted with the same participants for the first and second participations. The results of this study suggest that the score of each factor for PPD and bonding was slightly improved by parity in the first participations. However, PPD and bonding clearly showed moderate degree of relationships in the first

and second participations of the follow-up subgroup as well.

The changed scores of EPDS items and MIBS-J items by parity

Comparing the average scores of each EPDS and MIBS-J by parity, all items of the EPDS and MIBS-J decreased except for item 7 of the EPDS (Supplementary Table 2). Also, in the follow-up subgroup, the second participation scores of the same participants decreased compared with those of the first participation (Table 3).

In Japan, it became clear that the risk of PPD is significantly higher for primipara women than multiparous women (Mori et al., 2016; Morikawa et al., 2015; Takehara et al., 2018). The findings that the multiparous women showed better scores and the same participants improved in the second participation supported the results of previous studies. According to the factor analysis conducted based on the responses to the EPDS Japanese version, three factors were found; anhedonia from items 1 and 2; anxiety from items 3, 4, and 5; and depression from items 7, 8, and 9 (Kubota et al., 2014). In this study, the factors that helped improve the second participation scores with the same participants were anxiety and depression, and the AC scores were also improved in the MIBS-J. This might imply that Japanese people feel less anxiety and depression by going through the birth experience. Prior studies overseas report that multiparous women tend to fall into anxiety

and depression more frequently as compared to primiparas (Figueiredo and Conde, 2011), so improved PPD by parity might be associated with Japanese cultural factors (Morikawa et al., 2015).

The results of this study might indicate that Japanese primipara women are subject to suffer from anxiety and depression. Consequently, interventions for primiparas which would increase a child care experience would be a promising way to prevent or mitigate PPD and improve MIB.

In the future, we wish to perform additional follow-up studies over time after birth; however, in addition, other types of research such as intervention studies are needed to examine cause and effect relationships on this topic.

Limitation

Selective bias may have occurred in the follow-up subgroup. To be entered into the JECS study twice, the participants must have been pregnant during the recruitment period of 2011 to 2014, visited affiliated medical institutions for both pregnancies, and agreed to participate in the survey. It is possible that some participants experienced depressive symptoms with the first birth and did not consent to the survey with their second pregnancy. In addition, when the second survey was completed by the follow-up

subgroup, the responses of some participants might have been influenced by the memory of their first responses or social desirability (Kitamura et al., 1994; Windle, 1954, 1955). It is possible that scores might have been changed in EPDS and MIBS-J because of the influence of their memory of completing the survey from the first time.

Strength

The data used for the analyses came from a very large sample of women from all over Japan; therefore, our results could be represented the Japanese general population (Michikawa et al., 2018). We were also able to make comparisons between data for 2 consecutive births for the same women.

Conclusions

Our study revealed a robust, moderate relationship between PPD and MIB despite the influences of parity. However, many multiparous mothers' PPD and MIB scores decreased with the additional birth.

Acknowledgements

We are grateful to all of the participants of JECS and to all individuals involved in data collection. The Japan Environment and Children's Study was funded by the Ministry of the Environment, Japan. The findings and conclusions of this article are solely the responsibility of the authors and do not represent the official views of the above government agency.

Conflicts of interest statement

The authors declare no conflict of interest.

References

- Choi, H., Yamashita, T., Wada, Y., Narumoto, J., Nanri, H., Fujimori, A., Yamamoto, H., Nishizawa, S., Masaki, D., Fukui, K., 2010. Factors associated with postpartum depression and abusive behavior in mothers with infants. *Psychiatry Clin Neurosci* 64(2), 120-127.
- Cuijpers, P., Brannmark, J.G., van Straten, A., 2008. Psychological treatment of postpartum depression: a meta-analysis. *J Clin Psychol* 64(1), 103-118.
- Dennis, C.L., Hodnett, E., 2007. Psychosocial and psychological interventions for

treating postpartum depression. *Cochrane Database Syst Rev*(4), CD006116.

Dubber, S., Reck, C., Muller, M., Gawlik, S., 2015. Postpartum bonding: the role of perinatal depression, anxiety and maternal-fetal bonding during pregnancy. *Arch Womens Ment Health* 18(2), 187-195.

Edhborg, M., Nasreen, H.E., Kabir, Z.N., 2011. Impact of postpartum depressive and anxiety symptoms on mothers' emotional tie to their infants 2-3 months postpartum: a population-based study from rural Bangladesh. *Arch Womens Ment Health* 14(4), 307-316.

Figueiredo, B., Conde, A., 2011. Anxiety and depression symptoms in women and men from early pregnancy to 3-months postpartum: parity differences and effects. *J Affect Disord* 132(1-2), 146-157.

Figueiredo, B., Costa, R., Pacheco, A., Pais, A., 2009. Mother-to-infant emotional involvement at birth. *Matern Child Health J* 13(4), 539-549.

Hirokawa, K., Kimura, T., Ikehara, S., Honjo, K., Sato, T., Ueda, K., Iso, H., Japan, E., Children's Study, G., 2018. Associations between broader autism phenotype (BAP) and maternal attachment are moderated by maternal postpartum depression when infants are one month old: A prospective study of the Japan environment & children's study. *J Affect Disord* 243, 485-493.

Ishikawa, N., Goto, S., Murase, S., Kanai, A., Masuda, T., Aleksic, B., Usui, H., Ozaki, N., 2011. Prospective study of maternal depressive symptomatology among Japanese women. *J Psychosom Res* 71(4), 264-269.

Kawamoto, T., Nitta, H., Murata, K., Toda, E., Tsukamoto, N., Hasegawa, M., Yamagata, Z., Kayama, F., Kishi, R., Ohya, Y., Saito, H., Sago, H., Okuyama, M., Ogata, T., Yokoya, S., Koresawa, Y., Shibata, Y., Nakayama, S., Michikawa, T., Takeuchi, A., Satoh, H., Working Group of the Epidemiological Research for Children's Environmental, H., 2014. Rationale and study design of the Japan environment and children's study (JECS). *BMC Public Health* 14, 25.

Kitamura, T., Shima, S., Sugawara, M., Toda, M.A., 1994. Temporal variation of validity of self-rating questionnaires: repeated use of the General Health Questionnaire and Zung's Self-rating Depression Scale among women during antenatal and postnatal periods. *Acta Psychiatr Scand* 90(6), 446-450.

Kitamura, T., Takegata, M., Haruna, M., Yoshida, K., Yamashita, H., Murakami, M., Goto, Y., 2013. The Mother-Infant Bonding Scale: Factor Structure and Psychosocial Correlates of Parental Bonding Disorders in Japan. *Journal of Child and Family Studies* 24(2), 393-401.

Kitamura, T., Yoshida, K., Okano, T., Kinoshita, K., Hayashi, M., Toyoda, N., Ito, M.,

Kudo, N., Tada, K., Kanazawa, K., Sakumoto, K., Satoh, S., Furukawa, T., Nakano, H., 2006. Multicentre prospective study of perinatal depression in Japan: incidence and correlates of antenatal and postnatal depression. *Arch Womens Ment Health* 9(3), 121-130.

Kokubu, M., Okano, T., Sugiyama, T., 2012. Postnatal depression, maternal bonding failure, and negative attitudes towards pregnancy: a longitudinal study of pregnant women in Japan. *Arch Womens Ment Health* 15(3), 211-216.

Kubota, C., Okada, T., Aleksic, B., Nakamura, Y., Kunimoto, S., Morikawa, M., Shiino, T., Tamaji, A., Ohoka, H., Banno, N., Morita, T., Murase, S., Goto, S., Kanai, A., Masuda, T., Ando, M., Ozaki, N., 2014. Factor structure of the Japanese version of the Edinburgh Postnatal Depression Scale in the postpartum period. *PLoS One* 9(8), e103941.

Lefkovic, E., Baji, I., Rigo, J., 2014. Impact of maternal depression on pregnancies and on early attachment. *Infant Ment Health J* 35(4), 354-365.

Little, R.J.A., 1988. A Test of Missing Completely at Random for Multivariate Data with Missing Values. *Journal of the American Statistical Association* 83(404), 1198-1202.

Michikawa, T., Nitta, H., Nakayama, S.F., Yamazaki, S., Isobe, T., Tamura, K., Suda, E., Ono, M., Yonemoto, J., Iwai-Shimada, M., Kobayashi, Y., Suzuki, G., Kawamoto, T., 2018. Baseline Profile of Participants in the Japan Environment and Children's Study

(JECS). *J Epidemiol* 28(2), 99-104.

Mori, E., Maehara, K., Iwata, H., Tsuchiya, M., Sakajo, A., Ozawa, H., Aoki, K., Morita, A., Mochizuki, Y., Maekawa, T., 2016. Physical and psychosocial wellbeing of older primiparas during hospital stay after childbirth: A comparison of four groups by maternal age and parity. *Japanese Journal of Maternal Health* 56(4), 558-566.

Morikawa, M., Okada, T., Ando, M., Aleksic, B., Kunimoto, S., Nakamura, Y., Kubota, C., Uno, Y., Tamaji, A., Hayakawa, N., Furumura, K., Shiino, T., Morita, T., Ishikawa, N., Ohoka, H., Usui, H., Banno, N., Murase, S., Goto, S., Kanai, A., Masuda, T., Ozaki, N., 2015. Relationship between social support during pregnancy and postpartum depressive state: a prospective cohort study. *Sci Rep* 5, 10520.

Muzik, M., Bocknek, E.L., Broderick, A., Richardson, P., Rosenblum, K.L., Thelen, K., Seng, J.S., 2013. Mother-infant bonding impairment across the first 6 months postpartum: the primacy of psychopathology in women with childhood abuse and neglect histories. *Arch Womens Ment Health* 16(1), 29-38.

O'Higgins, M., Roberts, I.S., Glover, V., Taylor, A., 2013. Mother-child bonding at 1 year; associations with symptoms of postnatal depression and bonding in the first few weeks. *Arch Womens Ment Health* 16(5), 381-389.

Ohara, M., Nakatochi, M., Okada, T., Aleksic, B., Nakamura, Y., Shiino, T., Yamauchi,

A., Kubota, C., Morikawa, M., Murase, S., Goto, S., Kanai, A., Kato, R., Ando, M., Ozaki, N., 2018. Impact of perceived rearing and social support on bonding failure and depression among mothers: A longitudinal study of pregnant women. *J Psychiatr Res* 105, 71-77.

Ohara, M., Okada, T., Aleksic, B., Morikawa, M., Kubota, C., Nakamura, Y., Shiino, T., Yamauchi, A., Uno, Y., Murase, S., Goto, S., Kanai, A., Masuda, T., Nakatochi, M., Ando, M., Ozaki, N., 2017. Social support helps protect against perinatal bonding failure and depression among mothers: a prospective cohort study. *Sci Rep* 7(1), 9546.

Ohoka, H., Koide, T., Goto, S., Murase, S., Kanai, A., Masuda, T., Aleksic, B., Ishikawa, N., Furumura, K., Ozaki, N., 2014. Effects of maternal depressive symptomatology during pregnancy and the postpartum period on infant-mother attachment. *Psychiatry Clin Neurosci* 68(8), 631-639.

Okano, T., Masuji, F., Tamaki, R., Nomura, J., Murata, M., Miyaoka, H., Kitamura, T., 1996. Validation and reliability of Japanese version of the Edinburgh Postnatal Depression Scale (EPDS). *Arch. Psychiatri. Diagnos. Clin. Eval.* 7(4), 525-533.

Sockol, L.E., Battle, C.L., Howard, M., Davis, T., 2014. Correlates of impaired mother-infant bonding in a partial hospital program for perinatal women. *Arch Womens Ment Health* 17(5), 465-469.

Takehara, K., Tachibana, Y., Yoshida, K., Mori, R., Kakee, N., Kubo, T., 2018. Prevalence trends of pre- and postnatal depression in Japanese women: A population-based longitudinal study. *J Affect Disord* 225, 389-394.

Taylor, A., Atkins, R., Kumar, R., Adams, D., Glover, V., 2005. A new Mother-to-Infant Bonding Scale: links with early maternal mood. *Arch Womens Ment Health* 8(1), 45-51.

Tsuchid

a, A., Hamazaki, K., Matsumura, K., Miura, K., Kasamatsu, H., Inadera, H., et al (2019). Changes in the association between postpartum depression and mother-infant bonding by parity: Longitudinal results from the Japan Environment and Children's Study. *Journal of Psychiatric Research*, 110, 110-116.

van Bussel, J.C., Spitz, B., Demyttenaere, K., 2010. Three self-report questionnaires of the early mother-to-infant bond: reliability and validity of the Dutch version of the MPAS, PBQ and MIBS. *Arch Womens Ment Health* 13(5), 373-384.

van Buuren, S., 2007. Multiple imputation of discrete and continuous data by fully conditional specification. *Stat. Methods Med. Res.* 16(3), 219-242.

Windle, C., 1954. Test-Retest Effect on Personality Questionnaires. *Educational and Psychological Measurement* 14(4), 617-633.

Windle, C., 1955. Further Studies of Test-Retest Effect on Personality Questionnaires.

Educational and Psychological Measurement 15(3), 246-253.

Yalcin, S.S., Orun, E., Mutlu, B., Madendag, Y., Sinici, I., Dursun, A., Ozkara, H.A., Ustunyurt, Z., Kutluk, S., Yurdakok, K., 2010. Why are they having infant colic? A nested case-control study. *Paediatr Perinat Epidemiol* 24(6), 584-596.

Yamashita, H., Yoshida, K., Nakano, H., Tashiro, N., 2000. Postnatal depression in Japanese women. Detecting the early onset of postnatal depression by closely monitoring the postpartum mood. *J Affect Disord* 58(2), 145-154.

Yoshida, K., Yamashita, H., Conroy, S., Marks, M., Kumar, C., 2012. A Japanese version of Mother-to-Infant Bonding Scale: factor structure, longitudinal changes and links with maternal mood during the early postnatal period in Japanese mothers. *Arch Womens Ment Health* 15(5), 343-352.

Appendix

Members of the Japan Environment and Children's Study (JECS), as of 2017 (principal investigator, Toshihiro Kawamoto): Hirohisa Saito (National Center for Child Health and Development, Tokyo, Japan), Reiko Kishi (Hokkaido Regional Center for JECS, Hokkaido University, Sapporo, Japan), Nobuo Yaegashi (Miyagi Regional Center for JECS, Tohoku University, Sendai, Japan), Koichi Hashimoto (Fukushima Regional

Center for JECS, Fukushima Medical University, Fukushima, Japan), Chisato Mori (Chiba Regional Center for JECS, Chiba University, Chiba, Japan), Shuichi Ito (Kanagawa Regional Center for JECS, Yokohama City University, Yokohama, Japan), Zentaro Yamagata (Koshin Regional Center for JECS, University of Yamanashi, Chuo, Japan), Hidekuni Inadera (Toyama Regional Center for JECS, University of Toyama, Toyama, Japan), Michihiro Kamijima (Aichi Regional Center for JECS, Nagoya City University, Nagoya, Japan), Takeo Nakayama (Kyoto Regional Center for JECS, Kyoto University, Kyoto, Japan), Hiroyasu Iso (Osaka Regional Center for JECS, Osaka University, Suita, Japan), Masayuki Shima (Hyogo Regional Center for JECS, Hyogo College of Medicine, Nishinomiya, Japan), Yasuaki Hirooka (Tottori Regional Center for JECS, Tottori University, Yonago, Japan), Narufumi Suganuma (Kochi Regional Center for JECS, Kochi University, Nankoku, Japan), Koichi Kusuhara (Fukuoka Regional Center for JECS, University of Occupational and Environmental Health, Kitakyushu, Japan), and Takahiko Katoh (South Kyushu/Okinawa Regional Center for JECS, Kumamoto University, Kumamoto, Japan).

Supplementary Table 1 Demographic and obstetric characteristics for the follow-up subgroup

Characteristics	Subgroup		Subgroup	
	First participation		Second participation	
	(n = 3,753)		(n = 3,753)	
	Mean / n	(SD / %)	Mean / n	(SD / %)
Maternal age, mean (SD)	29.0	(4.4)	30.9	(4.5)
Postpartum depression, n (%)	541	(14.4)	383	(10.2)
Previous deliveries, n (%)				
Nullipara	2,437	(64.9)	3	(0.1)
Multipara	1,316	(35.1)	3,750	(99.9)
Highest educational level (year), n (%)				
<13	1,398	(37.3)	1,400	(37.3)
13-14	1,595	(42.5)	1,593	(42.5)
≥15	760	(20.3)	760	(20.3)
Annual household income (JPY), n (%)				
<4 million yen	1,661	(44.3)	1,590	(42.4)
4-5.9 million yen	1,195	(31.8)	1,267	(33.8)
≥6 million yen	897	(23.9)	896	(23.9)
Marital status, n (%)				
Married (including common law marriage)	3,559	(94.8)	3,736	(99.6)
Single (never married)	162	(4.3)	4	(0.1)
Divorced or widowed	32	(0.9)	13	(0.4)
Alcohol intake, n (%)				
Never drinker	3,367	(89.7)	3,413	(90.9)
Ex drinker	182	(4.9)	169	(4.5)
Current drinker	204	(5.4)	171	(4.6)
Smoking status, n (%)				
Never smoker	2,120	(56.5)	2,141	(57.1)
Ex smoker	1,444	(38.5)	1,431	(38.1)
Current smoker	189	(5.0)	181	(4.8)
Physical activity, yes (%)	2,752	(73.3)	2,589	(69.0)
History of anxiety disorder, yes (%)	89	(2.4)	87	(2.3)
History of depression, yes (%)	94	(2.5)	93	(2.5)
History of schizophrenia, yes (%)	5	(0.1)	2	(0.1)
Diagnostic record of mental disorder during pregnancy, yes (%)	25	(0.7)	19	(0.5)
Infant's anomaly, yes (%)	76	(2.0)	69	(1.8)
Infant's gender, n (%)				
Male	1,933	(51.5)	1,879	(50.1)
Female	1,820	(48.5)	1,874	(49.9)
Intensity and frequency of infant's crying, n (%)				
Quite often and long	648	(17.3)	409	(10.9)
Sometimes and short	2,925	(77.9)	3,132	(83.5)
Hardly	180	(4.8)	212	(5.7)

Supplementary Table 2. Means (SD) of EPDS score and bonding score in the cross-sectional group

Name of the scores or questions	Cross sectional group (N = 76,373)				
	Primipara (n = 32,342)		Multipara (n = 44,031)		*P
	mean	(SD)	mean	(SD)	
EPDS					
EPDS item 1	0.10	0.32	0.07	0.28	<.0001
EPDS item 2	0.12	0.36	0.09	0.32	<.0001
EPDS item 3	1.35	0.78	1.21	0.78	<.0001
EPDS item 4	1.31	0.85	1.00	0.82	<.0001
EPDS item 5	0.60	0.76	0.42	0.65	<.0001
EPDS item 6	1.31	0.60	1.20	0.57	<.0001
EPDS item 7	0.10	0.38	0.10	0.38	0.60
EPDS item 8	0.47	0.61	0.35	0.55	<.0001
EPDS item 9	0.13	0.39	0.11	0.35	<.0001
EPDS item 10	0.13	0.45	0.09	0.38	<.0001
MIBS-J					
MIBS-J item 1	0.07	0.27	0.06	0.25	<.0001
MIBS-J item 2	1.12	0.74	0.46	0.58	<.0001
MIBS-J item 3	0.07	0.28	0.06	0.26	<.0001
MIBS-J item 4	0.56	0.58	0.46	0.56	<.0001
MIBS-J item 5	0.11	0.34	0.05	0.24	<.0001