

衛生・生物化学 酵素生理学研究室

講師 赤尾光昭

◆ 原著

- 1) Heikal O.A., Miyashiro H., Akao T., and Hattori M. : Quantitative determination of paeoniflorin and its major metabolite, paeonimetabolin I, in the rat plasma by enzyme immunoassay. J. Trad. Med. 14: 15-19, 1997.
- 2) Kida H., Nakamura N., Meselphy M.R., Akao T., and Hattori M. : Isolation and identification of human intestinal bacteria capable of hydrolyzing saikosaponins. J. Trad. Med. 14: 34-40, 1997.
- 3) Heikal O.A., Akao T., Takeda S., and Hattori M. : Pharmacokinetic study of paeonimetabolin I, a major metabolite of paeoniflorin from peony roots. Biol. Pharm. Bull. 20: 517-521, 1997.
- 4) Heikal O.A., Kanaoka M., Akao T., and Hattori M. : Effects of spacer homologous and heterologous combinations on enzyme immunoassay for paeonimetabolin I, a major metabolite of paeoniflorin. J. Trad. Med. 14: 105-113, 1997.
- 5) Kida H., Akao T., Meselphy M.R., and Hattori M. : Enzymes responsible for the metabolism of saikosaponins from *Eubacterium* sp A-44, a human intestinal anaerobe. Biol. Pharm. Bull. 20: 1274-1278, 1997.

◆ 総説

- 1) Kobashi K., and Akao T. : Relation of intestinal bacteria to pharmacological effects of glycosides. Bioscience Microflora 16: 1-7, 1997.
- 2) Kobashi K., and Akao T. : Natural glycosides are prodrugs -Activation by human intestinal bacteria-. Bull. Soc. Pharmacognosy ROC No.V: 34-45, 1997.

◆ 学会報告

- 1) 来田浩明, 服部征雄, 赤尾光昭 : 柴胡サポニン代謝に関与するヒト腸内細菌由来の酵素について. 日本薬学会第117年会, 1997, 3, 東京.
- 2) Meselphy M.R., Heikal O.A., Akao T.,

Hattori M., Ono H., and Sadakane C. : Determination of paeoniflorin and paeonimetabolin I in rat plasma after oral administration of Toki-Shakuyaku-San (TS) and Shakuyaku-Kanzo-To (SK). 第13回和漢医薬学会大会, 1997, 8, 大阪.

◆ その他

- 1) Kobashi K., and Akao T. : Plant glycosides are natural prodrugs -The role of human intestinal flora, In Germfree Life and its Ramifications, by Hashimoto K. et al. (Ed.) 467-471, XII ISG Publishing Committee, Shiozawa, Japan, 1996.